

NUI MAYNOOTH
Ollscoil na hÉireann M^h Nuad

**Department of Geography
Rhetoric House
NUI Maynooth
Co Kildare**

**tel: ++ 353 (1) 7084566
fax: ++ 353 (1) 7083573
web: <http://icarus.nuim.ie>**

**EnviroCentre
10 Upper Crescent
Belfast
BT7 1NT**

**t 028 90278330
f 028 90278334
w www.envirocentre.co.uk
e belfast@envirocentre.co.uk**

Biodiversity and Climate Change in Ireland

Briefing Paper

John Coll¹
Cathy Maguire²
John Sweeney¹

**Submitted to Comhar SDC
January 2009**

¹ Irish Climate Analysis and Research Units, NUI Maynooth

² Envirocentre, Belfast

Table of Contents

1. Introduction	
1.1 Background.....	4
1.2 Rationale.....	5
1.3 Aims and objectives.....	6
2. Projected climate change impacts in Ireland	
2.1 Climate change impacts: terrestrial environment.....	8
2.2 Climate change impacts: aquatic environment.....	9
2.3 Biodiversity and ecosystem services.....	12
3. The Natura 2000 network in Ireland: coherence, connectivity and resilience	
3.1 Summary of the resource (Republic of Ireland)	14
3.2 Summary of the resource (Northern Ireland)	14
3.3 Structural connectivity: Natura 2000 network.....	17
2.4 Functional connectivity: terrestrial sites.....	18
2.5 Resilience: floodplains and river basin management	22
2.6 Resilience: coastal and estuarine settings.....	23
4. An initial assessment of habitats and species most at risk from climate change	
4.1 Habitats.....	25
4.2 Species.....	28
5. An overview of existing work.....	33
6. Development of policies for climate change and biodiversity	
6.1 Integration of climate change into biodiversity policies.....	35
6.2 Integration of biodiversity into climate change policies.....	35
6.3 Current mitigation and adaptation policy and impact on biodiversity.....	36
7. Gaps and sectoral policies	
7.1 Communication and knowledge gaps.....	37
7.2 Cross-sector synergies.....	38
7.2.1 Agriculture.....	38
7.2.2 Forestry.....	40
7.2.3 Water Management.....	41
7.2.4 Peatland resource management.....	41
7.2.5 Tourism.....	42
7.2.6 Other sectors.....	42
7.2.7 Gaps in policy making.....	42
8. Consider any emerging evidence of increased risk for the establishment of invasive alien species as a result of climate change.....	43

9. Capacity building and extending networks	
9.1 Biodiversity forum links.....	46
9.2 The case for a biodiversity research forum and suggested links.....	46
10. Developing an indicator species monitoring network	
10.1 Monitoring network: terrestrial settings.....	48
10.2 Monitoring network: marine settings.....	49
11. Concluding vision.....	51
12. Recommendations.....	54
Acknowledgements.....	55
Annex 1: Projected climate changes for Ireland.....	56
1.1 Climate model projections.....	56
1.2 Summary knowledge review: other drivers	
1.2.1 Sea level rise.....	60
1.2.2 Thermohaline circulation (THC) stability.....	60
1.2.3 Changes in storm and wave climate.....	61
Annex 2: An overview of the NPWs monitoring programme.....	63
References.....	66
Appendix 1: Glossary.....	79

1. Introduction

1.1 Background

The unprecedented social and economic changes experienced by Ireland over the past two decades has resulted in certain pressures on the environment increasing at a rate often exceeding that observed in other EU countries (EPA, 2008). Consequently, the four priority challenges identified by the EPA are:

- Limiting and adapting to climate change;
- reversing environmental degradation;
- mainstreaming environmental considerations;
- complying with environmental legislation and agreements (EPA, 2008).

Therefore the commissioning of a briefing paper on the impacts of climate change on biodiversity in Ireland is timely, not least since the Biodiversity Forum concludes that climate change is having and will continue to have a huge bearing on biodiversity in Ireland. The Biodiversity Forum also concludes that the maintenance of biodiversity has a role in ameliorating the impacts of a changing climate via the ecosystem services it provides. A knowledge review will also help inform the second National Biodiversity Plan (NBP) currently in development on the effects of climate change, invasive species and biosecurity, and in so doing will contribute to strategic national and local government policy planning over the next decade. With unprecedented levels of interest in biodiversity and the impacts of a changing climate, it has been recognised for some time that the state must take measures to develop or adapt existing national strategies and programmes for the conservation of biodiversity.

An obvious underpinning issue is that future climatic shifts could result in changes to species range dynamics which will reduce the relevance of present fixed protected areas for future conservation strategies. This is not to devalue the role our existing protected area networks have in the face of a changing climate, or to undermine their designation in protecting rare or threatened species (*sensu* J. Robinson and S. Egan pers. comm.). Rather it is a pragmatic recognition that these high quality and semi-natural natural areas will become important homes for new species and habitats and will remain important nodes in an overall site network (J. Robinson, pers. comm.).

Historically, conservation planning has mostly focused on preserving pattern and has acted reactively. However, the conservation agenda is now moving on to consider adaptation to climate change, and a landscape approach is more applicable for testing strategies such as habitat re-creation, and assessing ecosystem resilience (Thuiller *et al.*, 2008). There is also the need to assess and plan on the basis of functional connectivity rather than simple structural connectivity.

Implied in this is a further paradigm shift for conservation planning thinking. If unchecked climate change outstrips all possible available resources for new protected areas and as pathways for connectivity are exhausted, there may be a need to move away from the protected areas model in order to consider biodiversity conservation at the scale

of all potentially available suitable landscape units and habitats. Similarly, selecting new sites that offer a diverse range of physical habitats, even if they do not currently host rare or threatened species offers a way to sustain a wide diversity of species and constitutes a refreshed approach to site selection amid climate change uncertainties (J. Robinson, pers. comm.).

To illustrate the possible scale of the migration response for mobile species; recent work indicates an average 550km north-east range shift for European bird species in response to a 3° C end of century temperature increase (Huntley *et al.*, 2007); a rapid poleward movement along the continental shelf edge of ~ 1,100 Km for zooplankton species (Beaugrand *et al.*, 2002; Drinkwater *et al.*, 2003); northward shifts in the distribution of many commercially relevant fish species and sport fish species have also been documented (EEA, 2008; Irish Specimen Fish Committee, 2006, 2007). Consequently we consider that as well as providing a timely status audit as to how well 'climate proofed' Ireland's designated site network is, we propose that we can usefully extend the findings by examining how landscape-scale impacts to designated sites may best be modelled for Ireland in future work.

Predicted negative effects of climate change for Ireland's terrestrial and marine environments include changes in the distribution of species and the possible extinction of vulnerable species (EPA, 2008). Many aspects of Ireland's biodiversity are under threat, leading to habitat degradation and loss, and ultimately to an economic damage that still needs to be quantified (European Communities, 2008). The main threats arise from intensification of agriculture, poorly managed commercial forestry, peat extraction, land clearance and development, climate change and invasive alien species.

At a European level a recent European Environment Agency (EEA) report highlights that more action is needed towards halting biodiversity loss and maintaining the resilience of ecosystems because of their essential role in regulating the global climate system. Enhancing ecological coherence and the interconnectivity of the EU Natura 2000 network is key to the long-term survival of many species and habitats, for them to be able to adapt to a changing climate (EEA, 2008).

1.2 Rationale

Compelling evidence from around the globe indicates that species are already shifting their ranges in response to on-going changes in regional climates (Parmesan and Yohe, 2003; Root *et al.*, 2005; Walther *et al.*, 2005; Lavergne *et al.*, 2006), that species are altering their phenology (Menzel and Fabian, 1999; Visser and Holleman, 2001; White *et al.*, 2003; Zavaleta *et al.*, 2003; Jones *et al.*, 2006; Donnelly *et al.*, 2007) and that some species are facing extinction, or have become extinct (Parmesan, 2007; Pounds *et al.*, 2006; Foden *et al.*, 2007). Given the contemporary biodiversity crisis, effective conservation strategies that offset the climate change threats to species persistence will be critical in maintaining species and genetic diversity (Thuiller *et al.*, 2008).

These existing concerns are magnified since projected climate changes are likely to have an even greater impact on biota as the present century progresses (Berry *et al.*, 2002; Hill *et al.*, 2003; Thomas *et al.*, 2004; Thuiller *et al.*, 2005). Arising from these concerns there is a need in conservation planning to define at a landscape scale for any given region, the likely refugia corridors linking species' current and future ranges. In addition, conserving biodiversity as the climate changes is a two tiered challenge requiring adaptation (improved conservation strategies), together with mitigation.

There is considerable scientific consensus that the global climate is warming at a rate unprecedented in recent times and that warming trends are especially evident over northern hemisphere land areas and at high latitudes (Albritton *et al.*, 2001; Giorgi, 2005). In addition, numerous studies have shown that the most recent observations of changes in surface and free-atmosphere temperatures cannot be explained by (model-estimated) natural climate variability alone (Hegerl *et al.*, 2000; Barnett *et al.*, 2005).

However, there are significant remaining uncertainties in current predictions of future change at regionally and locally relevant scales. This is especially true for regions in North-western Europe and the North Atlantic, which are particularly challenging in terms of climate system understanding. Consequently planners are faced with a wide range of predicted changes from different models of unknown relative quality due to large, but unquantified, uncertainties in the modelling process (Cubasch *et al.*, 2001) (Annex Section 1.1). Nonetheless, and despite these issues surrounding the detail of outputs from different models, NW Europe continues to be identified as a regional climate change 'Hot-Spot' (*sensu* Giorgi, 2006).

Situated on the seaward western edge of north-western Europe and subject to both maritime and continental influences, the climate of Ireland is typified by spatial and temporal variability. These influences and the effects of the physical geography of the island produce a locally variable climate across the region. This variable climate contributes greatly to the biodiversity of the island, with a diverse mix of Atlantic, Arctic, Arctic-alpine and boreal elements occurring within a limited geographical area, and including many species on the edge of their global distributional range (Birks, 1997; Cross, 2006).

For example, the remaining woodlands of Ireland may be regarded as forming an extreme western and highly maritime extension of the west European transition or ecotone from Temperate Deciduous (Summer) Forest, through Boreal Coniferous Forest, to Boreal Deciduous Forest. Above the tree line and associated with the altitudinal decrease in temperature and increasing effects of wind, these communities are replaced by boreal alpine as well as sub-snowline and snowline associations where blanket bogs, heaths and dwarf shrub vegetation tend to be the dominants. It is this combination of highly variable local climates, together with many designated habitats of high conservation value being located in Ireland which necessitates the development of more locally relevant methods of climate change impact assessment for the biodiversity resource.

1.3 Aims and objectives

The aim of this briefing paper is to provide a review of available knowledge and the evidence base for policy on climate change and biodiversity. The specific objectives are:

- Provide an overview of the coherence, connectivity and resilience of National Parks, Nature Reserves, Natura 2000 sites and National Heritage Areas in order to maintain favourable conservation status of species and habitats in the face of the challenges presented by climate change.
- Make an initial assessment of the habitats and species in Ireland most at risk from climate change.
- Provide an overview of existing work which has been carried out on the potential impacts of climate change on biodiversity, including key habitats, and species.
- Assess synergies and gaps in the development of policies for climate change and biodiversity.
- Review climate change adaptation and mitigation policies and to report on the potential positive and negative impacts on biodiversity.
- Identify gaps and to make recommendations on how climate change considerations can be integrated into policy making in key sectors (including agriculture, forestry, inland waters, marine).
- Consider any emerging evidence of increased risk for the establishment of invasive alien species as a result of climate change.

2. Projected climate change impacts in Ireland

2.1 Climate change impacts: terrestrial environment

Superimposed on the biogeographical controls briefly outlined above, the landscapes of contemporary Ireland are the product of substantial human modification over a number of centuries, and ongoing human activities will be important in shaping the evolution of the landscape. The overall effect of these historical changes has been to convert a largely deciduous forested landscape, broken up by raised bogs in the lowlands and more extensive areas of blanket bog in the west and on the mountains, into a largely grassland-dominated landscape with small, scattered stands of native woodland and an increasing area of coniferous plantation (Cross, 2006).

Therefore in undertaking a review of climate change and biodiversity it is vital to account for other drivers of change. The impacts of predicted climate changes on habitats and species will be superimposed on other human drivers of change such as habitat fragmentation, agricultural change, eutrophication and invasive species. For example, the intensification and specialisation of agricultural practises have led to widespread declines in farmland bird diversity across much of Europe (McMahon *et al.*, 2008).

Therefore here, as with many other countries, the principal contemporary threat to biodiversity is habitat degradation and loss. This is especially true for a culturally-shaped landscape such as Ireland. The key policy drivers affecting the agricultural landscape are:

- Reform of the Common Agricultural Policy
- Implementation of the Water Framework Directive
- Nitrate Vulnerable Zones
- Renewable Energy Policy

Some drivers already operating may have increasing impacts with time (e.g. current grazing regimes, summer concentrations of ground-level ozone, or nitrogen deposition), new drivers may also emerge and an extreme climatic event may completely overwhelm any climate-change impacts. Legislation changes (driven by social, economic or environmental factors) could drive land management changes, such as reform of the Common Agricultural Policy causing a drastic decline in sheep numbers. Likewise the economics of land-use in Ireland could alter management, such as a decline in traditional land management practises, or an increase in land managed for conservation (Section 7.2.1). In the latter case, management may also be influenced by trends in management or advances in ecological knowledge, and by social awareness again leading to legislation changes. The changing patterns of land ownership and use across Ireland may result in a greater diversity of management aims, making the upland landscape more heterogeneous. Events or changes outwith Ireland may influence habitats and species, such as a change in Arctic breeding areas affecting over-wintering migratory birds.

It should be noted that these different drivers for change can interact in different ways and may combine with climate change in an additive or counteractive way. Indeed, such

unknown interactions may result in the most severe impacts. Given the potential for change arising from drivers such as those above, it is likely that such drivers may have greater impacts than climate change alone, certainly at least on a near term (2030s) horizon scan.

2.2 Climate change impacts: aquatic environment

Climate driven changes will also impact Ireland’s marine, coastal, littoral and freshwater communities, thereby affecting the wider biodiversity resource. The interactions between weather, ocean currents and sea temperature are complex: thus the detailed response of biological systems to changes is difficult to assess.

A wealth of marine habitats supports a wide diversity of life around Ireland’s coasts. This diversity is linked to the large scale movement of the North Atlantic Ocean’s thermohaline circulation (THC) (Annex Section 1.2.2). While the North Atlantic drift supplies warm waters on the west coast, cold sub-Arctic waters extend southwards to the Irish Sea to the east and juxtapose northern Boreal and southern Lusitanian marine species in a relatively small area.

Predicting changes in marine benthic habitats is difficult in Ireland due to the lack of data on species and habitat distributions, environmental requirements and understanding of how ecosystems function (Emblow *et al.*, 2003). In addition, no long-term datasets are available on species and habitats with which to examine patterns of marine environmental change over time (Emblow *et al.*, 2003). However, the MarClim study around UK coasts indicated that climate change is already having a profound impact on seashore indicator species (Laffoley *et al.*, 2005).

Harrison *et al.* (2001) examined the implications of climate change for the European marine environment, including five habitats of high conservation value which occur in Irish waters. They identified a range of direct and indirect effects which had to be considered (Table 1);

Table 1: Possible impacts on the marine environment considered by MONARCH (Harrison *et al.*, 2001) and other projects

Direct effects	Indirect effects
Sea-level rise	Change in thermohaline circulation*
Sea surface temperature change	Alteration in nutrient supply*
Increase in UV-B penetration*	Changes in wave climate*
	Changes in storminess*

*Insufficient evidence to predict likely scenarios

We have reviewed in summary form elsewhere (Annex Sections 1.2.1 – 1.2.3) evidence relating to prospective changes to some of these drivers. Given the present uncertainties surrounding the likely direction and magnitude of change for some of the key drivers, we conclude that as research continues and more sophisticated models continue to evolve, the scientific consensus is likely to change again over the coming decade. Therefore

there has to be an ongoing and iterative engagement between the policy and scientific research community (Section 6) to review biodiversity conservation planning on an ongoing basis.

In terms of the biological impacts of the present projected changes around Ireland's coasts, it is considered that there will be changes in the relative distributions between southern and northern species assemblages. Although the arrival of exotic and invasive species in coastal and estuarine habitats will have further implications, not least since the wider changes in climate may favour their increased establishment (Emblow *et al.*, 2003). However, see Emblow *et al.* (2003) for a fuller review and a discussion of caveats related to data and knowledge gaps. Table 7.9 (pp 178) (Emblow *et al.* 2003) also provides a useful summary. While Harrison *et al.* (2001, Chapter 6) provide further information on e.g. the possible impacts on estuarine birds around Irish and UK coasts which complements Emblow *et al.*'s (2003) assessment.

With warmer sea temperatures affecting phytoplankton communities, the resulting decline in sand eel populations would adversely affect a wide range of seabirds (Arkell *et al.*, 2007). Around the coasts of the UK e.g. evidence of another year of seabird breeding failure appears to be emerging (MCCIP, 2008). This most recent crash follows a series of repeated annual breeding failures which are increasingly considered to be climatically linked (Thompson and Ollason, 2001; Fredriksen *et al.*, 2004a; Grosbois and Thompson, 2005; Harris *et al.*, 2005)

There appears to be a strong link between warm winters, low sandeel biomass and poor breeding seabird performance (Rindorf *et al.*, 2000; Arnott and Ruxton, 2002; Wanless *et al.*, 2004, 2005). Sea warming appears to be disrupting the community structure and abundance of zooplankton on which hatching sandeel larvae may depend for survival, growth and recruitment (Fredriksen *et al.*, 2004b; Wanless *et al.*, 2004, 2005; Fredriksen *et al.*, 2006). Worryingly, the evidence suggests these repeated annual breeding failures linked to climate change impacts on their food sources are already substantially reducing populations of certain species (MCCIP, 2008).

In summary, associated with increased temperatures, there will be changes in the distribution of warm and cold water species in both coastal and offshore marine waters (Bolens *et al.*, 2005). Littoral and sub-littoral communities will be forced to respond to changing seasonal thermal regimes and light availability, a trend which will amplify over the century with increased warming. Irrespective of present climate change mitigation policies, the changes expected by the 2030s are already in train. Biodiversity is certain to be affected, although changes in community structure are perhaps more likely than changes in the numbers of species (Bolens *et al.*, 2005). To manage anticipated changes, there is increased impetus for an integrated coastal zone management system to be in place in order to better manage the drivers of change ahead of the 2030s.

Currently, the most challenging pollution issues relate to diffuse run off of pesticides and nutrient enrichment from agricultural land. With accelerated erosion triggered by a combination of exposure of the bare soil surface through human activity and extreme

rainfall, farming practises in coastal areas will have to account for the increased frequency of seasonal heavy rainfall events expected by the 2030s.

However, it is not only the future mobilization of pollutants from terrestrial sources which will become an issue for marine organisms, but the resuspension of historically deposited pollutants already in the sediments of coastal systems (especially estuaries) as seasonally rougher seas increasingly perturb the coastline. Increases reported over the past fifteen years in the concentration of dissolved organic carbon (DOC) in upland freshwaters elsewhere, are thought to be due to a combination of declining acid deposition and rising temperatures (e.g Evans *et al.*, 2005). The combination of higher summer temperatures and seasonal precipitation changes by the 2020s has the potential to increase carbon supply to lochs and estuaries, with any increase in turbidity of the water affecting levels of light and UV-B penetration. For example, Evans *et al.* (2005) report a step change in DOC mobilization from around 1996 for three of four monitored sites in Northern Ireland. While already, increased heavy rainfall events in recent summers are reported to be impacting ground-nesting birds and summer flooding of the Shannon Callows has impacted on Corncrake and other breeding waders (S. Egan, pers. comm.).

Perhaps more than for terrestrial settings, the direct impacts of climate driven change are likely to be more noticeable around the coasts and seas by the 2030s. It is not only the combination of localized changes to, for example, seasonal temperature and precipitation regimes that will drive change. Rather, prospective changes to wider hemispheric drivers such as the winter NAO, and its influence on water mixing characteristics and storm tracking will increasingly impact Irish marine and coastal communities. However, the reader is again referred back to some of the discussion in Annex Section 1.2. The effectiveness of managing realignment and dealing with pollutant mobilization in the coastal zone by the 2030s will largely depend on the extent to which present policy can integrate the necessary long term planning in a national coastal strategy which also integrates biodiversity considerations.

Seasonal rainfall changes will impact on upland catchments, particularly those in the west likely to exhibit rapid run-off rates. With any increase in heavy rainfall events will be an associated increase in sediment load and the delivery of suspended solids to fresh water and estuarine systems. In an analogous setting elsewhere, it has been reported that excessive stream discharges are dislodging fish spawning grounds in north-west Scotland (Kerr and Ellis, 2001). Similarly, it is thought that changes to flood frequencies are affecting communities of the listed freshwater pearl mussel (*Margaritifera margaritifera*) around NW Europe (Hastie *et al.* 2001, 2003).

Changes in the thermal environment will also directly impact upon some of Ireland's freshwater communities. In enclosed fresh water bodies, surface water temperatures will increase in response to higher summer temperatures. This will alter the mixing regime increasing thermal stratification (separation between surface and deeper waters which leads to deoxygenation) which will affect benthic and fish species. However, by the 2030s it is not so much mean seasonal changes which will drive such changes, but rather the increased likelihood of longer hot spells.

2.3 Climate change impacts: biodiversity and ecosystem services

In linking biodiversity and climate change issues, it is important to look beyond statutory obligations and notions of intrinsic worth. Biodiversity supports such diverse industries as agriculture, cosmetics, pharmaceuticals, pulp and paper, horticulture, construction and waste treatment. Consequently, the loss of biodiversity threatens our food supplies, opportunities for recreation and tourism, and sources of food, medicines and energy. However, there is a lack of knowledge regarding how ecosystems function and how biodiversity contributes to that function, across a range of scales, and it is generally considered that loss of biodiversity is likely to have a variety of effects on ecosystem function, resilience to change, the provision of goods and services and human well-being (Ferris, 2007).

In addition, the impact of losing elements of our biodiversity could deny future generations a wealth of cultural, scientific and commercial opportunities from new and emergent biotechnologies. Biodiversity is important for our health, for example as a source of pharmaceutical raw materials and also in terms of the quality of our food (Bullock *et al.*, 2008), while fully functioning ecosystems provide us with healthy and productive environments which support the economy of rural and coastal areas.

Following the momentum inspired by the *Stern Review of the Economics of Climate Change*, the G8+5 ministers expressed the need to explore a similar project on the economics of the loss of ecosystems and biodiversity (European Communities, 2008). Arising from the initial work there is already a call for national accounting systems to be more inclusive in order to measure the human welfare benefits that ecosystems and biodiversity provide (European Communities, 2008).

Other authors have suggested that we need to think afresh in terms of how e.g. policy for the Irish uplands can be formulated innovatively in the context of maximising provisioning, regulating and cultural services within an ecosystem services framework (Kirkpatrick, 2008). We suggest that exploring policy options integrating parallel ideas on how to implement management plans which merge strategic objectives in relation to restoration of habitats, future mitigation of climate change and amenity value may now be timely. An additional benefit is that the restoration of Ireland's upland landscape character could be a key component in ensuring that future revenue streams from the tourism sector are maintained, as well as extending the range and quality of ecosystem services provided.

Although only a preliminary and heavily conditioned figure, the current marginal value of ecosystems services in Ireland in terms of their contribution to productive output and human utility is estimated at over €2.6 billion per annum (Bullock *et al.*, 2008). While this estimate rests on only a few key examples and necessarily omits other significant services such as e.g. the waste assimilation by aquatic biodiversity and benefits to human health (Bullock *et al.*, 2008). Nonetheless, some provisional estimates of the economic contribution of biodiversity can be made across some sectors;

- Agriculture - ~ €1372 million per annum;
- Forestry - ~ €85 million per annum;
- Fisheries (catch) - ~ €180 million per annum;
- Aquaculture - ~€50 million per annum;
- Water quality - ~€385 million per annum;
- Human welfare - ~€330 million per annum (Bullock *et al.*, 2008).

By comparison the outlay spent directly on biodiversity protection is highly disproportionate to the value of the ecosystem services that biodiversity provides to the economy. For example, it is estimated that the National Parks and Wildlife Service spend ~€35 million per annum on protection, while the annual policy cost for the Rural Environment Protection Scheme (REPS) is ~€280 million.

To date no attempt has been made to attach a cost to some other significant and relevant ecosystem services related to biodiversity, such as:

- the buffering role of peatlands and other inland wetlands in retaining water during heavy rainfall events;
- the protective role of coastal wetlands in absorbing rising sea-levels and soft coastal defences in absorbing storm impacts;
- and the role of our peatlands, wetlands and woodlands in carbon capture and storage (S. Egan, pers. comm.).

Globally, tourism is one of the largest and fastest growing economic sectors and climate change is expected to have a range of direct (e.g. temperature rise) and indirect impacts. Wildlife tourism is growing in popularity and now generates substantial income for local economies around the world. For example, whale watching in Ireland was estimated to be worth €1,480,000 in direct revenues and €7,973,00 in indirect revenues in 1998 (Hoyt, 2000). While Ireland's recreational salmon fishery has been valued in excess of €100 million (CFB, 2000).

However, at regional and local scales there is little strategic planning in a locally based and highly fragmented sector due to a perception of climate change being a long-term global problem. At least part of Ireland's appeal for this growing sector is the richness and diversity of the habitats and species which contribute to the character of the landscape. There is therefore a need to assess the economic impact of possible biodiversity changes against new and emergent tourism markets of the future.

3. The Natura 2000 network in Ireland: coherence, connectivity and resilience

3.1 Summary of the resource (Republic of Ireland)

National Parks and Nature Reserves are areas designated for nature conservation and are in almost all cases entirely owned and managed by the National Parks and Wildlife Service (NPWS), constituting the most strictly protected conservation areas in the Republic of Ireland. There are currently 6 National Parks in the Republic of Ireland, which combined, cover 59,060 hectares (ha) (Craig 2001).

Natural Heritage Areas (NHAs) and Special Areas for Conservation (SACs), together with Special Protection Areas (SPAs) are designations that apply to areas of significant conservation value irrespective of ownership. The basic designation for wildlife is the NHA, defined as an area considered important for the habitats present or which holds species of plants and animals whose habitat needs protection (NPWS, 2008) (Figure 1).

To date, 75 raised bogs have been given legal protection, covering some 23,000 ha. These raised bogs are located mainly in the midlands; a further 73 blanket bogs, covering 37,000 ha, mostly in western areas are also designated as NHAs (NPWS, 2008) (Figure 1). There are also 630 proposed NHAs (pNHAs), which were published on a non-statutory basis in 1995, but have not since been statutorily proposed or designated.

The areas chosen as Special areas of Conservation (SAC) in Ireland cover an area of approximately 1, 337, 550 ha. Roughly 53% is land, the remainder being marine or large lakes and site areas range from ~0.01 ha to ~109, 000 ha. In addition, 121 Special Protection Areas (SPAs) have been designated since 1985, while 25 other sites are legally protected and will shortly be designated as SPAs. As yet, no marine SPAs exist despite a legal obligation to designate these. It should be noted that many existing and future SPAs overlap with SACs (Figure 2).

3.2 Summary of the resource (Northern Ireland)

While at present there are no national parks in Northern Ireland, there is a ministerial commitment to work towards the establishment of national parks, and in particular towards a Mourne National Park. In parallel with the work on primary legislation, specific work progressing the proposal for a Mourne National park is underway and a Mourne National Park Working Party is working up boundary proposals. As of 2005, the extent of designated sites in Northern Ireland comprised a total area of 93, 130 ha for Areas of Special Scientific Interest (ASSIs); 72, 549 ha for SPAs; 65, 943 ha for SACs and 77, 400 ha for Ramsar sites (EHS, 2008).

At the present time 52 Special Areas of Conservation have been designated and a further candidate SAC submitted to the European Commission (EC) as the major part of Northern Ireland's contribution to Natura 2000 (Figure 2). These sites have already been declared Areas of Special Scientific Interest (ASSIs). SPAs and SACs together form the European wide network of sites known as Natura 2000.




Figure 1: Outline map of Irish Natura 2000 site network. GIS shapefiles supplied by NPWS and the EPA (RoI) and the Department of the Environment (NI).




Figure 2: Outline map of Irish Natura 2000 network. Proposed NHAs (Republic of Ireland) overlain.

3.3 Structural connectivity: Natura 2000 network

It is clear from Figures 1 and 2 that there is considerable spatial overlap between sites within the various designations. This is particularly striking when the proposed NHAs for the Republic of Ireland (RoI) are overlain, particularly in relation to designated SACs (Figure 2). Another general feature is that there is more structural connectivity associated with western areas of the RoI, and to some extent around the coasts for both the RoI and Northern Ireland (NI). However, at the scale of the maps presented here, the areal extent of the designated sites reveals little of the detail of the landscape matrix and hence no real assessment of functional connectivity (Section 3.4).

In general, sites tend to reduce in spatial area and become more fragmented in the interior both sides of the border, this is not surprising as the designated areas are likely to be mirroring historical and contemporary patterns of settlement and land use. However, there is considerable structural connectivity associated with waterways and standing bodies of freshwater. This is likely to be significant when considering vectors which will facilitate the spread of aquatic invasive species associated with a changing climate (Section 8). Similarly, summer flooding of the Shannon Callows associated with any changes to extreme rainfall events may remain problematic for breeding birds given the impacts noted in recent summers.

Since the projected seasonal changes to mean seasonal temperature and precipitation regimes for Ireland are not of a high magnitude by the 2030s (Annex Section 1.1). It seems reasonable to consider that providing there are no dramatic policy, social, economic or environmental changes, and with climate change occurring as indicated in the scenarios; there will probably be no major noticeable effects on the structural connectivity of many of the designated terrestrial sites, with the possible exception of some of the smaller sites embedded within other areas of land use (Section 3.4).

However, currently degraded habitats may continue to degrade, such as through substrate erosion. Any major increases in current degradation, or new large-scale structural impacts will probably be attributable to alterations to management regimes brought about by other driving forces, such as CAP reform (Section 7.2.1).

Nonetheless, extreme events may have a very unpredictable and profound impact, depending on their frequency and intensity, and deserve further consideration. An extreme climatic event may completely overwhelm any climate-change impacts associated with mean changes, perhaps even changing the direction of succession or change in a very different way. For example, and given the structural connectivity associated with waterways noted above, flood events could have implications for biota at a catchment scale, such as for breeding birds along the Shannon Callows. Similarly, extreme events may be localized in time and space, such as storms causing bog-burst or landslips.

By contrast, extended drought may affect habitats over a wide spatial scale and last for several years. For example, the unusually hot and dry summer of 2003 had a huge impact on biomass productivity in Europe, with more than a year's recovery time having been estimated (Ciais *et al.*, 2005). Some authors have considered that this event was linked to

a multi-decadal pattern of variability in Atlantic Ocean forcing of western European summertime climate (Section 1.2.2). Even in the absence of other drivers and extreme events, and with no apparent change in many of the sites, there will still be a climate-induced momentum for change slowly building up in all communities that will exert increasing effects as the century progresses.

However, it is not adequate to consider the potential impacts of climate change on biodiversity in terms of structural changes to landscapes or habitats in isolation. There is also a need to consider how species and landscape interact, particularly in the intensively managed and fragmented landscapes of Ireland. Thus distinctions have to be made between the physical connectedness of habitats, and connectivity as a function of the ability of species to disperse through landscapes.

3.4 Functional connectivity: terrestrial sites

In order to halt the loss of biodiversity and meet other targets in the second Biodiversity Action Plan, there is a need to consider the impacts of climate change on species, both to understand their response and provision of potential adaptation measures. It is also vital to ensure that climate change adaptation and mitigation measures are not themselves harmful to biodiversity, and that increasingly an emphasis has to be placed on the importance of conserving biodiversity in the wider countryside, as well as in protected areas (EPA, 2008) (Section 1.1).

This is in addition to commitments in place which require reduction of the impacts of fragmentation, as well as further fragmentation that could potentially occur associated with rapid climate change. The recognition of behaviour as a link between process and pattern in landscape ecology is exemplified by the concept of functional connectivity, i.e. the degree to which the landscape facilitates or impedes movement among resource patches.

However, at the level of landscape planning the issue is further complicated since a contrasting impact of climate change is that invasive species may be able to spread further (Section 8). Therefore in order to protect native species assemblages, conservation interventions may be required to reduce connectivity for invasive species (Manchester and Bullock, 2000). Measures proposed to increase resilience in the face of biodiversity threats include expanded protected areas, varied and functional ecosystems and good habitat quality (Hopkins *et al.*, 2007; Mitchell *et al.*, 2007).

Consequently, when considering climate change adaptation measures the push and pull of these two conflicting management aims may best be considered in terms of re-building connectivity by way of engineering migration corridors in an already fragmented landscape. A key measure for increasing the speed at which species are able to respond to climate change is ensuring landscapes are permeable to species movement. Therefore there is a need to assess the dominant component of the landscape around and between protected areas.

The surrounding matrix has a significant impact on connectivity for many species in general. Semi-natural and extensive habitats are considered to be more conducive, or permeable, to species movement, while intensive land uses are regarded as less permeable, thereby reducing connectivity and effectively increasing ecological isolation (Murphy and Lovett-Doust, 2004; Watts *et al.*, 2005). However, while functional connectivity is related to structural connectivity, it has to be defined in relation to an individual species' requirements (Tischendorf and Fahrig, 2000; Hilty *et al.*, 2006; Taylor *et al.*, 2006). Thus for individual species, a landscape is functional if it allows a species to carry out all its ecological functions including movement for foraging, mate finding and dispersal. A basic principle of functional connectivity is that some land covers or land uses are more permeable than others (Adriaensen *et al.*, 2003; Donald and Evans, 2006).

While techniques have been developed to proactively target biodiversity conservation at a variety of spatial scales and which can use a simple structural approach or a more complex functional one (BEETLE; Watt *et al.*, 2005, 2007). Their application remains limited by:

- the availability of high quality spatial and species data; and
- restricted information on the dispersal and landscape permeability function for most species (Watt *et al.*, 2007).

This serves to highlight that while the concept of connectivity is relatively straightforward, translating and implementing it in the realm of practical conservation is not (Crooks and Sanjayan, 2006). A related issue is that while maintaining existing patches of semi-natural habitat in the landscape is important, creating new ones is desirable. However, the spatial locations of these patches in interaction with the biology of the species requiring to utilise them will largely determine their functional utility as 'stepping stones' for the suite of species requiring them (J. Robinson, pers. comm.).

With the appropriate species and habitat data available, it is clear that some of these methods could be adapted for climate change impact assessment for a number of Irish habitats and be applied to the protected area network when combined with outputs from climate models. Ideally these would also be combined with experiments to determine the actual impact of model predictions providing a more robust evidence base for policy. However, such research should proceed in tandem with initiatives which accelerate the implementation of known measures to make landscapes more permeable for species (S. Egan, pers. comm.) such as an expansion of existing REPS measures (Section 7.2.1).

For a maritime country like Ireland, it is also important that we consider the structural and functional linkages between habitats as extending from the lowlands to the uplands, therefore we also need to consider a vertical component in our adaptive landscape planning thinking. Associated with the typically steep climatic and hence environmental gradients of maritime uplands is an effective telescoping of broad latitudinal ecotones into a relatively small area. Therefore in considering the likely impacts of climate change, there is a need to maintain linkages between similar habitats in order to ensure

that species are able to track and colonise more favourable bioclimatic zones as these migrate upslope in a warming climate.

In summary, we need to develop a better understanding of spatial planning measures for Ireland which:

- implements the addition of known ‘stepping stone’ features such as wetlands and hedgerows which facilitate landscape permeability for a number of species;
- predict and integrate the impact of future weather patterns on priority habitats and species;
- review the effectiveness of implementation of landscape-scale adaptation initiatives; and
- link these to improved monitoring and recording programmes to facilitate model development.

One of the main shortcomings already identified in relation to managing Ireland’s biodiversity is the lack of data to provide baseline and up to date information on the distribution and abundance of certain species and habitats (EPA, 2008). However, this difficulty is not confined to the Republic as a similar situation also exists in Northern Ireland in relation to available baseline data for priority habitats and species (EHS, 2008). This lack of detailed high-resolution land cover and habitat data fit for national use hampers contemporary spatial planning measures relating to biodiversity, conservation and soil management (EPA, 2008). Issues like these contributed, at least in part, to a recent European review of Ireland’s biodiversity concluding that the conservation status of many of the most important habitats and species gave cause for concern (EPA, 2008; Figure 3). However, see also the NPWS report (2008) for qualifications surrounding the habitat assessments summarized in Figure 3.

It is clear therefore that if Ireland is to begin to address the challenges of linking biodiversity conservation policy to climate change mitigation and adaptation measures, these already significantly under-resourced areas (EPA, 2008) must be addressed. This review underscores and emphasizes these existing concerns, since if even the most basic and fundamental research is to be undertaken in relation to climate change impacts, access to high resolution spatial and species monitoring data is essential.

However, it should be emphasized that this is not a problem unique to Ireland. Species data can rarely be generated for all resolutions and all spatial extents, rather it tends to be available for large extents at coarse resolutions, or small extents at fine resolutions (Berry *et al.*, 2005). In addition, other criteria such as spatial representation, analytical soundness, measurability and potential inclusion in integrated assessment tools must also be considered (Erhard *et al.*, 2002).




Figure 3: Summary of Conservation status for all Annexed habitats (Source: NPWS, 2008)

Access to improved spatial data must also be accompanied by scaled-up research improving knowledge of topics such as species-area relationships and conceptual developments in the application of island biogeography theory; a better understanding of meta-population dynamics and plant and animal responses to landscape-scale structures and processes; and an improved understanding of abiotic processes within landscapes and the spatial extent of different controls.

Therefore as the century progresses, and even allowing for the uncertainties (Annex Section 1.1), key considerations for conservation of the biodiversity resource (in addition to those identified above) are:

- Expanding our protected areas and buffering these within larger habitat restoration schemes. Conserving Ireland’s peat bogs and restoring native woodland so they become both a useful future sequestration and conservation tool.
- These schemes would provide suitable conditions for existing populations, including space to move, as well as the space to accommodate new species in the future. Such schemes would contribute to continuing to provide vital ecosystem services such as carbon storage, water purification and flood management.
- The wider landscape must be permeable to species movements whether in an agricultural or urban context. Existing and future land-use policy should be reformed to secure and improve the site network in order to facilitate the delivery of a more permeable and resilient landscape.

3.5 Resilience: floodplains and river basin management

The level of vulnerability of wetlands to changes in climate depends on their position within hydrological landscapes. Hydrological landscapes are largely defined by the flow characteristics of ground water and surface water and by the interaction of atmospheric water, surface water, and ground water for any given locality or region. Assessment of these landscapes indicate that the vulnerability of all wetlands to climate change fall between two extremes: those dependent primarily on precipitation for their water supply are highly vulnerable, and those dependent primarily on discharge from regional ground water flow systems are the least vulnerable due to the greater buffering capacity of large ground water flow systems to climate change. Since most of the present water supply in Ireland comes from surface water, with between 20 and 25 percent supplied from groundwater (Charlton and Moore, 2003), Ireland's surface waters and the biological communities they support must therefore be considered as vulnerable to changes to seasonal precipitation regimes.

Wetland ecosystems are important because of the wide range of services which they perform, such as water regulation and purification (Millennium Ecosystem Assessment 2005) and carbon storage (Gorham 1991). Mitsch and Gosselink (1993) argue that hydrology is probably the most important determinant of the establishment and maintenance of specific wetland communities and wetland processes. Wetlands in Europe have already been identified as at risk from climate change through higher temperatures, greater evapotranspiration and altered precipitation amounts and patterns changing the hydrological regime (Hartig *et al.* 1997).

There is a general tendency for an enhanced seasonality of river flows in Ireland with projected climate changes:

- all areas will experience a decrease in summer runoff with a corresponding likelihood that the frequency and duration of low flows will increase in many areas;
- winter runoff is predicted to increase in the west, and particularly the north west, with a corresponding increase in the magnitude and frequency of flood events (Charlton and Moore, 2003; Charlton *et al.*, 2006).

These changes are linked to the projected changes to the spatial pattern of seasonal precipitation receipts across the country (Sweeney and Fealy, 2003, 2007).

Land use planning and land management and their effect on river flows are particularly critical. Improved management of floodplain meadows and other wetlands, removal of field drains, appropriate levels of grazing and improved woodland management in upland areas will all help to regulate water flow in a river catchment, and provide valuable habitats for wildlife. Therefore when considering the impacts of climate change it is also important to consider land use and human activity within the catchment, since these can have major impacts on the hydrological response to precipitation (Charlton and Moore, 2003; Charlton *et al.*, 2006).

In order to develop an appropriate management strategy to counteract climate change in streams, rivers and lakes, further consideration needs to be given to the impacts of (summer) low flow and more frequent flood events, as well as changes in water

temperature. A linked meta-model approach has recently been used to investigate the impacts of climate and socio-economic change scenarios on water flows and wetland species in two contrasting UK regions, but perhaps raised more questions than it did answers about the assumptions and limitations of the approach (Harrison *et al.*, 2008).

3.6 Resilience: coastal and estuarine settings

Coasts are projected to be exposed to increasing risks, including erosion, due to climate change and sea-level rise. The effect will be exacerbated by increasing human-induced pressures on coastal areas (IPCC, 2007). Approximately 300km² of land along Ireland's coastline is at risk of inundation from rising sea levels (McElwain and Sweeney, 2007). In Ireland, a higher platform of wave attack will inevitably mean greater erosion of soft coastlines. About 1metre of land retreat can be anticipated on sandy coastlines in Ireland for every centimetre rise in sea level (McElwain and Sweeney, 2007). Therefore if we incorporate the uncertainty range projections advocated by Hulme *et al.* (2002) as a first order approximation, this could translate to the loss of 4 -14 metres of soft coasts by the 2030s.

A rise in sea level effectively forces coastal habitats, such as sand dunes and mudflats, further in-land, although in some places the use of land and sea defences may prevent this landward migration of coastal habitats. The effective reduction in the area for the inter-tidal habitats often referred to as coastal squeeze affects species too. For example, migratory birds that rely on sand dunes and mudflats as staging and wintering posts may find their habitats reduced in area.

Around Ireland's coasts greater wave heights and storm surges may damage and remove sand dunes, machair, mudflats and shingle if these habitats are not allowed to migrate inland with changing coastal processes. These areas have a natural buffering effect, helping to absorb wave energy and may be viewed as natural protective strips around our coasts. However, the reader is also referred to some of the cautionary discussion in Annex Section 1.1 on climate model projections.

In coastal areas, as with all environments, there are subtle and cyclic interplays between physical processes and the biota exploiting the available habitats. Thus e.g. many soft coastlines undergo natural cycles of erosion and deposition linked to variations in sea level, sediment supply and wind and wave climate. These cycles may last from a few to many thousands of years and will be tracked by natural cycles of succession among colonising organisms. The obvious concern relating to climate change is that the rate and pace of changes will exceed the thresholds to which organisms are adapted.

Climate change is predicted to cause loss of inter-tidal habitat in low-lying areas e.g. coastal lagoons and estuaries (McElwain and Sweeney, 2007). Sand dune habitats and salt marshes are also vulnerable to sea-level and climate changes (IPCC, 2007). Recent modelling work in the UK suggests the most significant impacts are likely to be related to inter-tidal, saltmarsh and mudflat areas, which declined under all the sea-level rise scenarios considered (Gardiner *et al.*, 2007). Although the authors considered that compensation for these losses in some localities could be achieved through the creation of replacement

habitat by the managed realignment of sea defences, often in conjunction with engineered sediment supply to raise inter-tidal surfaces to levels conducive to vegetation establishment (Gardiner *et al.*, 2007).

For Ireland, many of the low lying estuarine sandflats, mudflats and lagoons found along the southeast coast, some of which have been identified as SACs, could be threatened (EPA, 2008). These habitats provide rich feeding grounds for a variety of bird species as well as important nursery grounds for juvenile fish. For a fuller review of the possible impacts on native and over-wintering birds around Ireland's coasts, the reader is directed to the work of Emblow *et al.* (2003).

While the setting back of sea defences to more landward locations (managed realignment) is a response to the loss of saltmarsh and mudflat habitats, in many locations new defence lines impinge upon coastal grazing marsh areas which are themselves often designated (Lee, 2001; Nicholls and Wilson, 2001; Pethick, 2002). By contrast, other low-lying coastal habitats, saltmarsh and estuaries which are prevented from extending landwards due to the presence of some fixed or artificial boundary area also at risk (EPA, 2008).

For these environments, recent work has indicated that management choices, which can be linked to socio-economic futures have a greater potential impact on habitat viability than climate change. Therefore the choices that society makes will be key to the protection and conservation of biodiversity in the coastal zone (Richards *et al.*, 2008).

Consequently, there is a further need to identify and develop the tools that make it possible to manage Ireland's wetland and coastal ecosystems in the face of climate change. In addition, mechanisms are needed which allow climate change considerations to be integrated with management of human activities, identification of conservation sites, and the processes for monitoring, assessing and reporting on the status of habitats, species and ecosystems.

Further assessments are needed on the likely impact of the combined effects of ocean acidification and climate change, with the development of a composite index to track change. In summary, further research is required in a number of key areas including:

- Identifying constraints and opportunities and providing recommendations for optimal adaptation to climate change in coastal zones.
- Investigating the implications of climate change for the biodiversity of coastal wetlands.
- Improving understanding of the biodiversity impacts of managed retreat practices.
- Evaluating the biodiversity implications of building hard defences.
- Assessing the implications of ocean acidification associated with climate change in relation to the impact on ecosystem function and the biodiversity of Irish coastal communities.

4. An initial assessment of habitats and species most at risk from climate change

While we have touched upon a number of related areas in the settings review above, here we intend to focus on habitats and species considered particularly vulnerable or at risk from climate change and for which assessments are available from the literature. We present these alongside an existing status review for Annexed habitats and species in order to provide an indication of climate change vulnerability alongside contemporary status audits.

4.1 Habitats

Below we provide a summary table taken from the NPWS (2008) assessment with an added column providing an indication of climate change impact knowledge status. We define vulnerability as the extent to which a natural or social system is susceptible to sustaining damage from climate change (IPCC, 2007). The footnotes beneath the table summarise the information sources.

Table 2: Assessment of each attribute and overall Conservation Status for Annexed Habitats (* indicates priority habitat). Definitions of the non-climate change terms are supplied in NPWS, 2008.

EU Code	Habitat Names (summarised)	Range	Area	Structure & Functions (Condition)	Future Prospects	Overall	Climate Change Vulnerability Assessment
1110	Sandbanks	Good	Good	Good	Poor	Poor	Unknown
1130	Estuaries	Good	Good	Unknown	Poor	Poor	Unknown
1140	Tidal Mudflats and Sandflats	Good	Good	Poor	Poor	Poor	Unknown
1150	Coastal Lagoons*	Good	Poor	Bad	Poor	Bad	Unknown
1160	Large Shallow Inlets and Bays	Good	Good	Unknown	Poor	Poor	Unknown
1170	Reefs	Good	Unknown	Poor	Poor	Poor	Unknown
1210	Annual Vegetation of Drift Lines	Good	Poor	Good	Poor	Poor	Unknown
1220	Perennial Vegetation of Stony Banks	Good	Poor	Poor	Poor	Poor	Unknown
1230	Vegetated Sea Cliffs	Good	Good	Poor	Poor	Poor	Unknown ¹
1310	Salicornia mud	Good	Poor	Poor	Poor	Poor	Unknown
1320	<i>Spartina</i> Swards	Good	Poor	Good	Poor	Poor	Unknown
1330	Atlantic Salt Meadows	Good	Poor	Poor	Poor	Poor	Unknown
1410	Mediterranean Salt Meadows	Good	Good	Poor	Poor	Poor	Unknown
1420	Halophilous Scrub	Good	Bad	Poor	Bad	Bad	Unknown
2110	Embryonic Shifting Dunes	Good	Poor	Poor	Poor	Poor	Unknown
2120	Marram Dunes (White Dunes)	Good	Bad	Bad	Bad	Bad	Favoured ¹
2130	Fixed Dunes (Grey Dunes)*	Good	Poor	Bad	Bad	Bad	Favoured ¹
2140	Decalcified Empetrum Dunes*	Good	Good	Bad	Poor	Bad	Unknown
2150	Decalcified Dune Heath*	Good	Good	Bad	Poor	Bad	Unknown
2170	Dunes with Creeping Willow	Good	Good	Poor	Poor	Poor	Unknown
2190	Humid Dune Slacks	Good	Poor	Poor	Bad	Bad	Unknown
2IAO	Machair*	Good	Poor	Bad	Bad	Bad	Low ¹
3110	Lowland Oligotrophic Lakes	Good	Good	Bad	Bad	Bad	Unknown
3130	Upland Oligotrophic Lakes	Good	Good	Bad	Bad	Bad	Unknown

3140	Hard Water Lakes	Good	Good	Bad	Bad	Bad	Unknown
3150	Natural Eutrophic Lakes	Unknown	Unknown	Unknown	Bad	Bad	Unknown
3160	Dystrophic Lakes	Good	Unknown	Bad	Bad	Bad	Unknown
3180	Turloughs*	Good	Good	Poor	Poor	Poor	Medium-High ¹
3260	Floating River Vegetation	Good	Good	Bad	Bad	Bad	Unknown
3270	Chenopodium rubri	Good	Good	Good	Good	Good	Unknown
4010	Wet Heath	Good	Unknown	Bad	Bad	Bad	Medium ¹
4030	Dry Heath	Good	Good	Poor	Poor	Poor	Low ¹
4060	Alpine and Subalpine Heath	Good	Poor	Poor	Poor	Poor	Medium-High ¹
5130	Juniper Scrub	Good	Poor	Poor	Poor	Poor	Unknown
6130	Calaminarian Grassland	Good	Good	Good	Poor	Poor	High ¹
6210	Orchid-Rich Grassland/Calcareous Grassland*	Good	Bad	Bad	Bad	Bad	Unknown
6230	Species-Rich Nardus Upland Grassland*	Good	Bad	Bad	Bad	Bad	Unknown
6410	Molinia Meadows	Good	Bad	Bad	Bad	Bad	Unknown
6430	Hydrophilous Tall Herb	Good	Good	Poor	Poor	Poor	Unknown
6510	Lowland Hay Meadows	Bad	Bad	Bad	Bad	Bad	Low ¹
7110	Raised Bog (Active)*	Bad	Bad	Bad	Bad	Bad	Medium-High ¹
7120	Degraded Raised Bogs	Good	Good	Poor	Poor	Poor	Medium-High ¹
7130	Blanket Bog (Active)*	Good	Bad	Poor	Bad	Bad	Medium-High ¹
7140	Transition Mires	Good	Good	Bad	Bad	Bad	Unknown
7150	Rhynchosporion Depressions	Good	Good	Good	Good	Good	Unknown
7210	Cladium Fens*	Good	Good	Bad	Bad	Bad	Medium-High ¹
7220	Petrifying Springs*	Good	Good	Bad	Bad	Bad	Medium-High ¹
7230	Alkaline Fens	Good	Good	Bad	Bad	Bad	Medium-High ¹
8110	Siliceous Scree	Good	Poor	Poor	Poor	Poor	Unknown ¹
8120	Calcareous Scree	Good	Poor	Poor	Poor	Poor	Unknown
8210	Calcareous Rocky Slopes	Good	Poor	Poor	Poor	Poor	Unknown
8220	Siliceous Rocky Slopes	Good	Poor	Poor	Poor	Poor	Unknown
8240	Limestone Pavement*	Good	Poor	Poor	Poor	Poor	Unknown ¹
8310	Caves	Good	Unknown	Good	Good	Good	Unknown
8330	Sea Caves	Good	Unknown	Good	Good	Good	Unknown
91A0	Old Oak Woodlands	Good	Bad	Bad	Bad	Bad	Medium ¹
91D0	Bog Woodland*	Good	Poor	Poor	Poor	Poor	Medium-High ¹
91E0	Residual Alluvial Forests*	Good	Bad	Bad	Bad	Bad	Unknown
91J0	Yew Woodlands*	Bad	Bad	Bad	Bad	Bad	Unknown

1 – Byrne *et al.* (2003)

For the interested reader seeking further information, the habitat distribution maps (NPWS, 2008) are available at:

<http://www.npws.ie/en/PublicationsLiterature/HabitatsDirectiveReport07/Habitats/>

The approach is useful in identifying that climate change impacts may be superimposed on an already poor outlook for a number of priority habitats. Based on this initial qualitative assessment, the most vulnerable habitats overall appear to be;

- 7110 Raised Bog (Active);
- 7130 Blanket Bog (Active);
- 7210 Cladium Fens;
- 7220 Petrifying Springs;

- 3180 Turloughs;
- 91D0 Bog Woodland.

The exercise also indicates that priority habitats requiring further climate change impacts research appear to be;

- 1150 Coastal Lagoons;
- 2140 Decalcified Empetrum Dunes;
- 2150 Decalcified Dune Heath;
- 6210 Orchid-Rich Grassland/Calcareous Grassland;
- 6230 Species-Rich Nardus Upland Grassland;
- 91E0 Residual Alluvial Forests;
- 91J0 Yew Woodlands.

The key contemporary threats to these important habitats and species have been identified as direct habitat damage; overgrazing and undergrazing; water pollution; unsustainable exploitation; invasive alien species and recreational pressures (NPWS, 2008). In addition, in January 2007 the European court of Justice ruled that Ireland did not have in place a system of strict protection for Annex IV species (Table 3) due to inadequate monitoring arrangements and deficient procedures to protect species from development projects (C-183/05) (Clenaghan, 2008). Certain derogations under the Wildlife Act, 1976 were also found to be incompatible with EU law (Clenaghan, 2008).

Similar driving forces and pressures over the past century have also seen a decline in numbers and the loss of species in priority habitats in Northern Ireland (EHS, 2008). Consequently, thirteen priority habitats including lowland meadow, species rich hedgerows, blanket bog and coastal sand dunes are considered to be in decline (EHS, 2008). Similarly, and mirroring the situation in the Republic, information on the conservation status and trends is not currently available for many habitats in Northern Ireland due to inadequate monitoring, although plans are in place to improve the monitoring programme (EHS, 2008).

4.2 Species

Again we provide a summary table taken from the NPWS (2008) assessment with an added column providing an indication of climate change impact knowledge status. We define vulnerability as above, and the footnotes beneath the table again summarise the information sources.

Table 3: Assessment of each attribute and overall Conservation Status for Annexed Species. Definitions of the non-climate change terms are supplied in NPWS, 2008.

EU Code	Species Name	Annex	Range	Population	Suitable Habitat	Future Prospects	Overall	Climate Change Vulnerability Assessment
1421	Killarney Fern (<i>Trichomanes speciosum</i>)	II, IV	Good	Good	Good	Good	Good	Medium ^{1,2}
1528	Marsh Saxifrage (<i>Saxifraga hirculus</i>)	II, IV	Good	Good	Good	Good	Good	Unknown ^{1,2}
1833	Slender Naiad (<i>Najas flexilis</i>)	II, IV	Good	Poor	Poor	Good	Poor	Medium ^{1,2}
1393	Slender Green Feather-Moss (<i>Hamatocaulis vernicosus</i>)	II	Good	Good	Good	Good	Good	Unknown
1395	Petalwort (<i>Petalophyllum ralfsii</i>)	II	Good	Good	Good	Good	Good	Unknown
1376	Maerl (<i>Lithothamnion corralloides</i>)	V	Good	Unknown	Unknown	Poor	Poor	Favoured ³
1377	Maerl (<i>Phymatolithon calcareum</i>)	V	Good	Unknown	Unknown	Poor	Poor	Unknown
1400	White Cushion Moss (<i>Leucobryum glaucum</i>)	V	Good	Good	Poor	Good	Poor	Unknown
1409	<i>Sphagnum</i> genus	V	Good	Good	Poor	Poor	Poor	Unknown
1413	<i>Lycopodium</i> species group	V	Good	Poor	Poor	Poor	Poor	Medium ²
5113	<i>Cladonia</i> subgenus <i>Cladina</i>	V	Good	Good	Poor	Poor	Poor	Unknown
1013	Geyer's Whorl Snail (<i>Vertigo geyeri</i>)	II	Good	Poor	Poor	Poor	Poor	Unknown
1014	Narrow-mouthed Whorl Snail (<i>Vertigo angustior</i>)	II	Good	Poor	Poor	Poor	Poor	Low ¹
1016	Desmoulin's Whorl Snail (<i>Vertigo moulinsiana</i>)	II	Bad	Bad	Poor	Bad	Bad	Unknown
1024	Kerry Slug (<i>Geomalacus maculosus</i>)	II, IV	Good	Good	Good	Good	Good	Favoured ¹
1029	Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>)	II, V	Good	Bad	Bad	Bad	Bad	High ⁴
1990	Nore Freshwater Pearl Mussel (<i>Margaritifera durrovensis</i>)	II, V	Bad	Bad	Bad	Bad	Bad	High ⁴
1092	White-Clawed Crayfish (<i>Austropotamobius pallipes</i>)	II, V	Poor	Poor	Poor	Poor	Poor	Unknown
1065	Marsh Fritillary (<i>Euphydryas aurinia</i>)	II	Good	Poor	Poor	Poor	Poor	Unknown
1095	Sea Lamprey (<i>Petromyzon marinus</i>)	II	Poor	Poor	Poor	Poor	Poor	Unknown
1099	River Lamprey (<i>Lampetra fluviatilis</i>)	II, V	Good	Good	Good	Good	Good	Unknown
1096	Brook Lamprey (<i>Lampetra planeri</i>)	II	Good	Good	Good	Good	Good	Unknown
1102	Allis Shad (<i>Alosa alosa</i>)	II, V	Good	Unknown	Unknown	Unknown	Unknown	Medium ⁵
5046	Killarney Shad (<i>Alosa fallax killarnensis</i>)	II, V	Good	Good	Good	Good	Good	Medium ⁵
1103	Twaite Shad (<i>Alosa fallax fallax</i>)	II, V	Good	Bad	Unknown	Poor	Bad	Medium ⁵
5076	Pollan (<i>Coregonus autumnalis</i>)	V	Good	Bad	Poor	Poor	Bad	High ⁵
1106	Atlantic Salmon (<i>Salmo salar</i>)	II, V	Good	Bad	Poor	Poor	Bad	Medium-High ¹
1202	Natterjack Toad (<i>Bufo calamita</i>)	IV	Bad	Bad	Poor	Poor	Bad	Low-Medium ¹
1213	Common Frog (<i>Rana temporaria</i>)	V	Good	Good	Poor	Good	Poor	Unknown
1223	Leatherback Turtle (<i>Dermochelys coriacea</i>)	IV	Unknown	Unknown	Unknown	Poor	Poor	Unknown
1303	Lesser Horseshoe Bat (<i>Rhinolophus hipposideros</i>)	II, IV	Good	Good	Good	Good	Good	Low ¹
1309	Common Pipistrelle (<i>Pipistrellus pipistrellus</i>)	IV	Good	Good	Good	Good	Good	Unknown
5009	Soprano Pipistrelle (<i>Pipistrellus pygmaeus</i>)	IV	Good	Good	Good	Good	Good	Unknown
1317	Nathusius' Pipistrelle (<i>Pipistrellus nathusii</i>)	IV	Good	Good	Good	Good	Good	Unknown

1322	Natterer's Bat (<i>Myotis nattereri</i>)	IV	Good	Good	Good	Good	Good	Unknown
1314	Daubenton's Bat (<i>Myotis daubentonii</i>)	IV	Good	Good	Good	Good	Good	Unknown
1330	Whiskered Bat (<i>Myotis mystacinus</i>)	IV	Good	Good	Good	Good	Good	Unknown
1320	Brandt's Bat (<i>Myotis brandtii</i>)	IV	Good	Good	Good	Good	Good	Unknown
1326	Brown Long-Eared Bat (<i>Plecotus auritus</i>)	IV	Good	Good	Good	Good	Good	Unknown
1331	Leisler's Bat (<i>Nyctalus leisleri</i>)	IV	Good	Good	Good	Good	Good	Unknown
1334	Irish Hare (<i>Lepus timidus hibernicus</i>)	V	Good	Unknown	Poor	Good	Poor	Unknown
1355	Otter (<i>Lutra lutra</i>)	II, IV	Good	Poor	Good	Good	Poor	Unknown
1357	Pine Marten (<i>Martes martes</i>)	V	Good	Good	Good	Good	Good	Low-Medium ¹
1364	Grey Seal (<i>Halichoerus grypus</i>)	II, V	Unknown	Good	Good	Good	Good	Unknown
1365	Common (Harbour) Seal (<i>Phoca vitulina vitulina</i>)	II, V	Unknown	Good	Good	Good	Good	Unknown
1345	Humpback Whale (<i>Megaptera novaeangliae</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
1349	Bottle-Nosed Dolphin (<i>Tursiops truncatus</i>)	II, IV	Good	Unknown	Good	Good	Good	Unknown
1350	Common Dolphin (<i>Delphinus delphis</i>)	IV	Good	Unknown	Good	Good	Good	Unknown
1351	Harbour Porpoise (<i>Phocoena phocoena</i>)	II, IV	Good	Good	Good	Good	Good	Unknown
2027	Killer Whale (<i>Orcinus orca</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2029	Long-Finned Pilot Whale (<i>Globicephala melas</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2030	Risso's Dolphin (<i>Grampus griseus</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2031	White-Sided Dolphin (<i>Lagenorhynchus acutus</i>)	IV	Good	Unknown	Good	Good	Good	Unknown
2032	White-Beaked Dolphin (<i>Lagenorhynchus albirostris</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2034	Striped Dolphin (<i>Stenella coeruleoalba</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2035	Cuvier's Beaked Whale (<i>Ziphius cavirostris</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2038	Sowerby's Beaked Whale (<i>Mesoplodon bidens</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2618	Minke Whale (<i>Balaenoptera acutorostrata</i>)	IV	Good	Unknown	Good	Good	Good	Unknown
2621	Fin Whale (<i>Balaenoptera physalus</i>)	IV	Good	Unknown	Good	Good	Good	Unknown
5020	Blue Whale (<i>Balaenoptera musculus</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
5031	Sperm Whale (<i>Physeter catodon</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
5033	Northern Bottlenose Whale (<i>Hyperoodon ampullatus</i>)	IV	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
2619	Sei Whale (<i>Balaenoptera borealis</i>)	IV	Unknown	Unknown	Good	Unknown	Unknown	Unknown
1348	Northern Right Whale (<i>Eubalaena glacialis</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown
5029	False Killer Whale (<i>Delphinapterus lucas</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown
2037	True's Beaked Whale (<i>Mesoplodon mirus</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown
2622	Pygmy Sperm Whale (<i>Kogia breviceps</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown
5029	Beluga/White Whale (<i>Delphinapterus leucas</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown
5034	Gervais' Beaked Whale (<i>Mesoplodon europaeus</i>)	IV	Vagrant	Vagrant	Vagrant	Vagrant	Vagrant	Unknown

1 – Byrne *et al.* (2003)

2 – Wyse-Jackson (2008)

3 – Harrison *et al.* (2001)

4 – Hastie *et al.* (2001, 2003)

5 – FSBI (2007)

It should be noted that the Byrne *et al.* (2003) assessments are based on a later century (2050s and 2080s) horizon scan than we provide here. However, for consistency we apply their vulnerability ascription. While for the Marsh Saxifrage (*Saxifraga hirculus*) there are conflicting interpretations.

For the interested reader seeking further information, species distribution maps (NPWS, 2008) are available at:

<http://www.npws.ie/en/PublicationsLiterature/HabitatsDirectiveReport07/Species/>

There are similar knowledge gaps in relation to the identified 271 Northern Ireland priority species. Hence the lack of datasets or baseline information for many species prevents reporting trends and flags the need for targeted systematic monitoring of Northern Ireland priority species (EHS, 2008). Although not listed here, all the Annex I species of the Birds Directive could have been included here as these inform the designated SPAs which comprise the other half of the Natura 2000 network. Similarly, the impacts of climate change alongside other drivers has been identified as a threat to bird species on the All Ireland red and Amber lists (Lynas *et al.*, 2007).

Considering the poor assessment for a number of our freshwater species, and given that the plight of the Freshwater Pearl Mussel has been previously highlighted at Ministerial level (NPWS, 2008). We see considerable potential for linkages to initiatives aimed at implementing the Water Framework Directive (WFD) (Section 7.2.3). Not least since a primary environmental objective of the WFD for surface water is that the ecological and chemical status of all water bodies will be good or high by 2015, and that in no case will the status deteriorate below its present condition (EPA, 2008). Therefore in contrast to existing Water Directives seeking to protect specific uses of water, the WFD is concerned, *inter alia*, with the protection of aquatic ecosystems as a whole (EPA, 2008) (Figure 4).




Figure 4: A summary schematic of the scope of all waters the WFD aims to protect (Source: www.wfdvisual.com)

In particular, we see a useful concordance of timing in the planning cycles for allowing the integration of biodiversity and climate change issues into the WFD programme. Since the production of River Basin Management Plans (RBMPs) on a planning cycle commencing in 2009 and running every six years thereafter are at the core of the new participatory management assessment and monitoring programmes. We see considerable utility for the integration of species-specific climate change considerations for threatened species at an appropriate spatial scale and on a similar timescale.

In addition, and since each RBMP cycle is a continuous and iterative process of assessment, review and action to ensure the final RBMP meets the aims and objectives of the WFD (EPA, 2008). We identify a further and useful congruence with the decision support frameworks we identify in relation to biodiversity and climate change considerations in Section 7.1, including the iterative framework we identify in Figure 5.

We also see a congruent timescale for the 2030s horizon scan we provide here for biodiversity. Thus e.g., key milestones for the implementation of the WFD include:

- 2009 – Finalise river basin management plan including programme of measures and begin first management cycle;

- 2012 – Make operational programmes of measures;
- 2015 – Meet environmental objectives; first management cycle ends; second river basin management plan and first flood risk management plan;
- 2021 – Second management cycle ends;
- 2027 – Third management cycle ends, final deadline for meeting objectives (EPA, 2008).

5. An overview of existing work

Below we provide a summary table providing an overview of work which has been done in relation to biodiversity and climate change in Ireland. The interested reader is directed to the reference lists of the above for further information, as well as to the recent state of knowledge review by Clenaghan (2008) and the references therein.

Table 4: Summary of literature, biodiversity and climate change

Year	Author	Type	Notes
2001	Harrison <i>et al.</i> (MONARCH)	Report section	Impacts on species and habitats, UK and Ireland
2003	Byrne <i>et al.</i>	Report section	Climate change and biodiversity assessment
2003	Emblow <i>et al.</i>	Report section	Climate change and the marine environment
2004	Donnelly <i>et al.</i>	Journal paper	Climate change indicators for Ireland
2005	Berry <i>et al.</i> (MONARCH)	Report section	Impacts on species and habitats, UK and Ireland
2006	Jones <i>et al.</i>	Journal paper	Vegetation response to climate change
2007	Donnelly <i>et al.</i>	Report section	Climate change & changing phenologies in Ireland. Impacts on semi-natural ecosystems.
2007	Walmsley <i>et al.</i> (MONARCH)	Report section	Impacts on species, UK and Ireland
2007	Arnell <i>et al.</i>	Report	Cross-sector scoping, Northern Ireland
2007	Iremonger <i>et al.</i>	Report	Forest biodiversity
2007	Huntley <i>et al.</i>	Book	Modelled range shifts, European bird species
2008	Wyse-Jackson, P.	Report	List of climate change vulnerable plant species
2008	FSBI	Briefing paper	Climate change and the fishes of Britain and Ireland

6. Development of policies for climate change and biodiversity

In European policy a temperature increase of 2°C has emerged as the threshold for acceptable climate change and the benchmark temperature against which to consider atmospheric concentrations of greenhouse gases and emission reduction profiles. A 2°C rise in temperature equates to stabilisation at 450 ppm carbon dioxide equivalent (CO₂e) (Anderson and Bows, 2008). Climate change policy is informed by this 2°C threshold even though stabilising at 450 ppm CO₂e offers only a 46% chance of not exceeding 2°C (Meinshausen, 2006). Recent research by Anderson and Bows (2008) which examined the difference between empirical and modelled data on emissions since 2000 indicates that that given current emission trends (3% annual emission growth), 2°C may provide a reasonable guide for mitigation, but it is a misleading basis for informing adaptation policy which would be much better guided by stabilisation at 650 ppm CO₂e, equating to approximately a 4°C rise in temperature.

This review is taking place at a time when policy both in the Republic of Ireland and Northern Ireland is in development. In both jurisdictions the current policy framework does not take an integrated approach to climate and biodiversity. In the Republic of Ireland, key strategies such as the Biodiversity Strategy and the National Adaptation Plan are in revision and development respectively and are not likely to be forthcoming until next year.

In Northern Ireland, the policy framework is also in development and will be shaped by the Climate Change Bill which has just recently been passed by the UK parliament and will require the development of an adaptation programme. The Northern Ireland Climate Change Impacts Partnership (NICCIP) has been established to widen the understanding and knowledge of the impacts of climate change and necessary adaptation actions within Northern Ireland. Key objectives are to promote ownership across sectors and increase adaptation capacity. One objective of the research project on preparing for a changing climate in Northern Ireland (Arkell *et al.*, 2007) was to produce an adaptation strategy and identify the public sector bodies responsible for delivery. The research highlighted the lack of sector specific risk assessments for Northern Ireland and concluded that as a result awareness, willingness to change and a general sense of urgency to consider adaptation was lacking. The report also highlighted how research on impacts has not been carried through into policy and there has been no clear coordination of strategic planning within and between the bodies responsible to address climate change risks and adaptation. Last month the NI Assembly Environment Committee after hearing evidence from the Tyndall Centre based on the analysis in Anderson and Bows (2008) announced an inquiry into climate change, so there is some uncertainty over Northern Ireland policy and whether targets from the Climate Change Bill will be adopted. In light of the Climate Change Bill and assembly inquiry, it is likely that there will be further development of the devolved policy agenda in the coming months.

In considering current policy and guidance, it is necessary to examine both the integration of climate change considerations into biodiversity policy and the integration of biodiversity considerations into climate change policy. The timing of the review

coincides with an absence of key policies and strategies which makes it difficult to assess their effectiveness and what the impact will be on biodiversity. However, this lack of an integrated policy framework in combination with the timetable for policy development provides an opportunity for the synergies between these areas to be fully realized and reflected in a joined up policy framework.

6.1 Integration of climate change into biodiversity policies

The integration of climate change into biodiversity strategies and policies is proceeding and the revised Biodiversity Strategy will have climate change as a key theme. Many of the actions contained in the strategy will be focused at integrating biodiversity considerations into other policy areas and processes and this provides an opportunity to put in place actions integrating climate change considerations as well.

There are a range of other policies and plans focused on protecting and managing the biodiversity resource and these can also integrate climate change. Habitat action plans and species action plans can and do integrate climate change as a pressure and/or threat and contain management actions. Site management plans such as SAC conservation management plans also offer potential to integrate climate change more fully. At present these have a five year time frame (2005-2010) and integrating climate change considerations fully will require a longer time perspective and this could be addressed in the next round of plan development with specific consideration of the potential impact of climate change and the inclusion of mitigation actions into management plans and policies. The implementation of policy to produce management plans for SPAs would also be a useful complimentary measure if climate change adaptation measures could be integrated. In addition as highlighted in Section 4.2, we see the WFD as an important vehicle for integrating climate change and biodiversity policies for the aquatic environment

Public awareness campaigns on biodiversity have also been undertaken in both Northern Ireland (It's In Our Nature) and the Republic of Ireland (Notice Nature). Notice Nature targeted sectors with production of guidelines for construction, business and tourism. Although these do not specifically mention climate change and the need to factor in climate change considerations when planning to protect biodiversity. Given the level of uncertainty over specific impacts this may have been too complex an issue for these focused guidelines to include. However climate change should be incorporated more explicitly into any future phases of public awareness campaigns. Notice Nature also targeted the agricultural sector although the main provision for addressing biodiversity issues was through REPS.

6.2 Integration of biodiversity into climate change policies

There has been little to no integration of biodiversity issues into climate change policy to date in Ireland, however the National Climate Change Strategy recognizes the impacts of climate change on biodiversity. The National Climate Change Strategy also contains a commitment to produce a National Adaptation Strategy by 2009. This is in the very early stages of development and a steering group has now been established. The National Adaptation Strategy provides both an opportunity and a vehicle for integrating

biodiversity and climate change issues and could direct the consideration of both into a wide range of strategies and plans including development plans and sector-specific guidance. For biodiversity issues there is a record of cooperation and collaboration on an island wide basis, it would also be beneficial to develop adaptation policies relating to biodiversity on a biogeographical basis.

6.3 Current mitigation and adaptation policy and positive and negative impacts on biodiversity

As adaptation policy is currently in development an assessment of the positive and negative impacts on biodiversity is difficult to do. Current mitigation actions including the use of REPS and biomass cultivation could have either positive or negative effects depending on how the policies are developed and implemented. There is also the EU policy dimension and certain sectors such as agriculture, fisheries and energy are largely integrated at an EU level through the single market and common policies and a recent green paper from the European Commission (EC, 2007) identifies a range of options for EU actions on adaptation.

However there are some current mitigation policies that are planned that could potentially have a negative impact on biodiversity unless the proper evaluations are undertaken. Specific actions include:

- The need for a Strategic Environmental Assessment (SEA) of renewable energy policy.
- Evaluation of REPS measures for their impacts on biodiversity and effectiveness.
- Assessment of changes in land use to ensure they are not impacting negatively on biodiversity.
- Sustainability Impact Assessments (SIAs) should be carried out to ensure biodiversity is taken into account in the development of renewable energy resources.
- Encouragement of biofuel crops that do not impact on biodiversity.

7. Gaps and sectoral policies

7.1 Communication and knowledge gaps

It has been recognized for a number of years that a clear policy framework is a prerequisite for the delivery of critical and coherent decisions relating to biodiversity policy (NPBR, 2003). There is also an urgent need to ensure that environmental policy making in general and biodiversity policy making in particular are informed by well-founded scientific knowledge (NPBR, 2003). However, as we have reviewed in preceding sections, difficult adaptation and mitigation decisions relating to projected changes in climate have to be taken despite significant scientific uncertainties. Consequently, as in a number of other sectors, there are substantial gaps between science and practice in the area of climate change and biodiversity.

A related problem arises from the innate tension between policy maker's expectations and the ability of science to deliver the outcomes required by policy, this is especially true in relation to biodiversity adaptation to climate change. These sorts of tensions can be attributed to;

- (i) The acquisition rate of ecological (research) knowledge is slow relative to climate change. Consequently policy makers have to implement decisions on adaptation despite imperfect information.
- (ii) Science and policy-maker communication also plays an important role. Thus while high level communication exists with respect to broad themes (e.g. the conservation of biodiversity during climate change), such broad themes may not be easily broken down into questions that can be readily tackled by scientific research (Sutherland *et al.*, 2006).

However, there is also a tension for the scientific community since the reductionist paradigm of classical science is not best suited to addressing the complexity inherent in conducting climate change impact assessments (CCIAs), but rather a systems-based approach linked across disciplines is required. Benefits arising from improved communications between the communities could include:

- more realistic expectations from policy makers on what science can deliver and a better definition of questions that can be addressed by scientists;
- a better recognition on the part of researchers of what constitutes genuine policy needs; and
- a recognition that applied research can also equate to good science.

Given the wider uncertainties and some of these innate tensions, an iterative engagement between science and policy which is both reflexive and outcome centered is envisaged as being the most appropriate way to proceed (Figure 5). Importantly, such a risk-averse decision making framework allows for a regular re-iteration of the problem and an ongoing re-evaluation of the objectives in light of any new or emerging scientific knowledge.




Figure 5: A framework to support good decision making in the face of climate change risk (Willows and Connell, 2003).

Aside from these very real communication issues, we also identify a further need to examine the cross-sectoral synergies of climate change and how these will impact upon biodiversity. Not least since despite the significant amount of climate related research in this sector, there is an apparent lack of co-ordination of policy development and strategic planning between the various responsible bodies to address climate change risks and adaptation (Arkell *et al.*, 2007). Specifically, we see related climate change driven impacts in how the agricultural, forestry, water management and tourism sectors are managed as being particularly important in relation to future biodiversity conservation.

7.2 Cross-sector synergies

Below we identify a number of key sectors where we consider it essential that biodiversity and climate change considerations should be integrated into existing policy. We then briefly explore some of what we consider to be some of the key issues and likely drivers of change.

7.2.1 Agriculture

As the principal land use, farming has important linkages with the tourism and leisure industries, as well as having the dominant influence on the visual characteristics of Ireland's landscape. The total land area of Ireland is approximately 6.9 million hectares of which 4.3 million hectares or 62% is used for agriculture.

One major effect of the CAP Reform on the Irish landscape is likely to be a change in grazing systems. The current predominance of sheep grazing is largely due to subsidies received from the EU, as opposed to market forces. The introduction of Single Farm

Payments, decoupling and removal of subsidies will likely result in a significant reduction in sheep numbers and a proportional increase in cattle grazing. Large changes in grazing such as these may alter the sward mix of grasslands on extensive systems as cattle are less discriminating grazers. There may also be some interaction with climate change impacts since wetter soils may be more vulnerable to erosion and compaction from cattle poaching under climate change. However, this is dependant upon the breed of, with evidence that some traditional breeds had a lesser impact on soil resources (H. Kirkpatrick, pers comm.).

A further effect of decoupling could be that due to market forces growers may choose to produce non-food crops. Oilseed rape, for example has numerous industrial applications, and large-scale expansion in land used for this purpose will alter the visual characteristics of the landscape. Cross Compliance and Good Agricultural and Environmental Condition (GAEC) requirements (as part of the CAP reform) will also instigate some changes in the visual landscape. These may be seen through changes in land management such as controlled grazing of cattle or re-forestation.

The implementation of the Water Framework Directive and Nitrate Vulnerable Zones will undoubtedly change the soil and water environment in Irish farmland. The two major implications for farming in Ireland will be the control of diffuse pollution from agricultural sources and regulatory control of abstraction for irrigation purposes. These have the potential to alter the underlying characteristics of farmland through changes in the purity and volume of water in rivers and streams. Restrictions from these policy mechanisms on timing of water abstraction and application of fertilisers may conflict with agricultural needs under an altered climate.

The substitution of fossil fuels with the increasing use of renewable energy sources is now recognised by governments worldwide as a fundamental priority in reducing greenhouse gas emissions. The main sources of renewable energy will be wind power, biomass, hydro, tidal and wave energy. As a consequence the visual characteristics of the agricultural landscape could alter quite dramatically as land is used for afforestation and short rotation coppice for renewable purposes.

Similarly, if government policy encourages the uptake of bio-energy; miscanthus and oilseed rape (used for bio-diesel) could radically alter the visual characteristics of the farming landscape, particularly if they replace grassland production. Climate change would improve the yields of some energy crops in Ireland. These issues were recognised in a recent Conference of the Parties (COP, 2008) call to Governments and others to continue to investigate and monitor the positive and negative impacts of the production of biofuels on biodiversity.

As energy crops are still a relatively new venture in Ireland, the Department of Agriculture and Food is introducing measures to stimulate production at farm level at an estimated cost of €14 million over the period 2007-2009 (CAP, 2006). The measures include a new national payment of €80 per hectare to stimulate production of energy crops. At present, farmers can avail of an EU premium of €45 per hectare under the EU Energy Crops Scheme, to grow energy crops intended for use in the production of biofuels and biomass (CAP, 2006). Approximately 2,000 hectares was claimed under this in 2006. As an additional incentive, the new €80 national payment will be paid as a top-

up to the existing €45 EU premium and increases the overall premium available to €125 per hectare (CAP, 2006).

It is clear that any or all of the above changes could have major implications for biodiversity. An obvious positive measure would be an expansion of REPS measures dealing with hedgerows, habitats, field margins, and biodiversity options such as the provision of nature corridors, species-rich grassland, tree planting and the environmental management of set-aside. While carbon sinks will be created primarily by Ireland's afforestation measures, these could be complemented significantly by support through REPS for the creation and maintenance of hedgerows e.g. In addition, the expansion of many of these existing REPS measures would help address a number of the connectivity issues in relation to climate change adaptation we have identified previously.

7.2.2 Forestry

The Republic of Ireland published a strategic plan for the forest sector in 1996 (DAFF, 1996) which involved increasing the forest cover dramatically. Ireland is one of the least forested countries in Europe, even though forestry plantations have increased forest cover from less than 1% of land cover to about 10% in the last century. The plan aims to increase this to 17% by 2030, mainly by planting new commercial forests at approximately 20,000 ha per year (Iremonger *et al.*, 2007). This significant increase represents a large change in land use and land cover across Ireland, and has far reaching economic, social and ecological consequences.

As Ireland's forest authority, the Forest Service has a clear strategy for conserving and enhancing biodiversity in forests. The role of the Forest Service is to ensure that forestry practice in Ireland conforms to the principles of sustainable forest management (SFM), whereby forestry develops in a way that maximises its contribution to national economic and social well-being on a sustainable basis and is compatible with the protection of the environment (McAree, 2002). The Forestry Inspectorate is responsible for ensuring that all conditions relating to biodiversity are complied with in grant-aided forests, and it plays a pivotal role in the policing, monitoring and promotion of these important requirements (McAree, 2002).

However, in order to promote forest biodiversity and fully practice SFM, it is necessary to know what organisms are associated with the forest plantations, and what the manager should be aiming at. The various projects undertaken by Iremonger *et al.* (2007) concluded that forestry plantations can make a significant positive contribution to biodiversity in the landscape if properly planned and managed, and can have a negative effect if not. Another conclusion was that the promotion of biodiversity in forestry needs the support of good policies and practices (Iremonger *et al.*, 2007).

The Planning and Management Tools for Biodiversity in a Range of Irish Forests (PLANFORBIO) research programme supports targeted forest biodiversity research in the contemporary Irish landscape (UCC, 2008). In keeping with government strategy to increase planting of forests that support a wealth of biodiversity throughout Ireland ongoing fieldwork as part of the wider project is concerned largely with mixed species plantations, while continuing to survey our native woodlands (UCC, 2008).

Obviously these sorts of programmes could be integrated as part of a wider collaboration examining what climate change could mean for biodiversity in Ireland's forests. We broach some of the strategic benefits such linkages could bring further in Section 9.2. In addition, and providing mixed species plantations comprise much of the new woodland, this would help address some of the connectivity issues explored in Sections 3.3 and 3.4, as well as the direct sequestration of atmospheric carbon dioxide.

7.2.3 Water management

The Water Framework Directive (WFD) 2000/60/EC has significant interconnections and linkages with other EU legislation. These include:

- Environmental Assessment Directives; Environmental Impact Assessment (EIA) - 85/337/EEC as amended by 97/11/EC and 2003/35/EC;
- and Strategic Environmental Assessment (SEA) – 2001/42/EC;
- Public Participation Directive (2003/35/EC), and the Birds (79/409/EEC) and Habitats (92/43/EC) Directives (Bennet and Sheate, 2007).

It is clear that data issues remain uppermost in terms of the challenges faced by River Basin Districts (RBDs) and conservation bodies implementing the Birds and Habitats Directives. Baseline data from different processes, particularly in relation to conservation, need to be improved as a matter of urgency both in terms of quality of data gathered and its availability (Bennet and Sheate, 2007). We would consider that this is an imperative given the bleak outlook identified for some of our freshwater species in Table 3 above.

7.2.4 Peatland resource management

Given the long historical and ongoing contemporary interventions involved in shaping the landscapes of Ireland, and the subsequent gradations between landscape categories. It is perhaps somewhat artificial to consider bog and peatland habitats as distinctive landscape types. For example, extensive areas of peatland forest exist and their management into the future is an important issue in a climate change context as conifer species may spread or possibly become more invasive in a changing climate (C. Douglas, pers. comm.). This has considerable implications for the biodiversity reservoir associated with Ireland's organic soil resource.

There are contemporary concerns about the siting of windfarms on bog and heath sites, as well as related concerns surrounding the targeting of afforested peatland for windfarm developments (C. Douglas, pers. comm.). These relate to a number of peat slides apparently being associated with windfarm developments in various parts of the country. Associated with the loss of soil cover are changes in the hydrology and continued functioning in relation to biodiversity, as well as carbon storage and sequestration. Therefore as we increasingly drive towards renewable technologies to meet our emissions targets, there is a need to link site selection criteria with biodiversity considerations.

Also, and given that the functioning of our peatlands are intrinsically linked to their hydrology. We can see clear linkages and synergies with the RBMPs of the WFD in

terms of integrating climate change adaptation measures for the biodiversity resource of our peatlands.

7.2.5 Tourism

We have briefly explored some of the issues here previously (Section 1.3) and conclude that wildlife tourism is a sector which is likely to continue to grow in the future. However, preservation of Ireland's landscape character and the restoration of degraded habitats are a key component in ensuring that future revenue streams from the tourism sector are maintained. Similarly, we envisage that restoration of landscape connectivity could be part of a win-win scenario in the context of wider biodiversity conservation, given the connectivity and buffering issues we explored in Sections 3.3 and 3.4. We also conclude that further research is needed into how patterns of tourism may change with a changing climate and what the implications of this are for biodiversity.

7.2.6 Other sectors

The growing impact on biodiversity of invasive species means it is necessary to also consider how those sectors which introduce invasive species and are therefore key to reducing the threat, such as horticulture, aquaculture, the pet and aquaria trade and construction are considering climate change in policy making. The Invasive Species Ireland project which is jointly funded by NPWS and the Northern Ireland Environment Agency (NIEA) has started to produce Codes of Practice for these key sectors in combination with a programme of stakeholder engagement. Further incorporation of climate change considerations into the outputs and next phase of this project could be an effective way of raising awareness, ensuring legislative compliance and getting actions taken by these sectors.

7.2.7 Gaps in policy making

A review of policy and guidance in key sectors has identified a lack of integrated policy on climate change and biodiversity. Some specific gaps include:

- Strategic Environmental Assessment of energy policy.
- Lack of guidance on integration of biodiversity and climate considerations into planning policy both at national and local level.
- Lack of guidelines for local authorities and other sectors on biodiversity and climate.
- Lack of biodiversity and climate change considerations in public procurement policies and practices.

There are a number of ways that these gaps can be addressed. One is the full implementation of the Strategic Environmental Assessment Directive which will strengthen biodiversity and climate considerations in sectoral policies. Another is the integration of climate proofing into Environmental Impact Assessment and ensuring that policy impact assessments address impacts on ecosystems supported by instruments that internalise the costs of damages to natural capital and ecosystem services (EC, 2007). However the key policy driver for addressing climate change and biodiversity could be achieved by ensuring that the National Adaptation Plan is integrated with and harmonized with the Biodiversity Strategy and that harmonisation of adaptation policy also takes place on an all-island basis.

8. Evidence for increased risk from invasive species as a result of climate change

There is a general scientific consensus that climate change is likely to favour invasive species, leading to new invasions and spread of already established invasive species. A recent review on climate change and invasive species prepared for the standing committee of the Bern Convention concluded that there is increasing evidence that climate change will affect the processes underlying biological invasions but that the current state of knowledge is not robust enough to make specific predictions (Capdevila and Zilletti, 2008). As highlighted in other sections of this review, ecosystems will differ in their responses to elevated CO₂ and temperatures and this is also the same for invasive species. Predictions need to be made on a species by species basis taking into account the biology and ecology of the species, the susceptibility of the habitat to invasion, the vulnerability of native biodiversity to climate change and the interactions between ecosystems and human activities.

Invasive species can also exacerbate the impacts of climate change by impacting biodiversity in advance of direct climatic impacts. Climate change will interact with invasive species through different mechanisms such as changes in pathways and vectors, habitat alteration increasing vulnerability to invasion and changes in range distribution of potential and established invaders. Climate change will also influence propagule pressure, that is, the number and frequency of invasive species arriving in a particular habitat. Increases in mean temperature are also likely to lead to northward expansion of populations.

In terrestrial ecosystems climate change may affect the dynamics of invasions by causing changes in native ecosystems that increase their vulnerability to invasion and by favouring traits of specific invasive species. Temperature and length of growing season are key variables determining the distribution of invasive plant species. Changes in these could influence the reproductive capacity of species and even their reproductive strategies. In general models reveal that simulated climate change negative impacts on native ecosystems are likely to facilitate invasions (Thuiller *et al.*, 2007). However it is imperative that experiments are also undertaken to evaluate invasive species responses to changes in variables such as elevated CO₂, temperature and precipitation regime to provide an evidence base for policy and decision making.

In the aquatic environment, climate change will affect many ecological processes and dispersal pathways and these will affect the success of invasive species. For marine distributions, extreme meteorological events may be as important as gradual climate changes. Extremes or sudden changes in temperature, salinity and turbidity may favour some exotic species. Changes to sea level, in combination with storm events, are likely to result in alterations to low lying areas and concomitant changes to water temperature, circulation, retention and sediment redistribution are likely to modify native species assemblages and could provide niches for exotics (Emblow *et al.*, 2003). Recently there has been recruitment of the non-native Pacific oyster (*Crassostrea gigas*) in response to

elevated summer temperatures. This species was introduced for aquaculture and water temperatures were considered to low for successful breeding. Recruitment has been documented in a number of sea lochs and research is now ongoing into the potential impact.

The Invasive Species Ireland (ISI) project which began in May 2006 aims to reduce the impact and threats from invasive species on the island of Ireland and is a joint initiative between the National Parks and Wildlife Service and the Northern Ireland Environment Agency. In Ireland, practical management of invasive species is challenging because of the cross-border implications of controlling introductions and spread. A pro-active stance is fundamental since prevention of introductions is demonstrably more cost-effective than reactive control or eradication measures. The ISI project has put in place practical steps that aim to minimise introductions, evaluated management and control measures, and laid the foundations for a cross-jurisdictional framework to respond to species introductions.

A key part of the project is to undertake a review of progress after two years and explore options for a way forward. Climate change was identified as a key issue that needs to be incorporated into the next phase of ISI, some specific actions that need to be undertaken include the revision of the risk assessments to incorporate climate change to identify species that will present a growing threat to native biodiversity in the context of climate change.

Invasive Species Ireland has carried out over 600 risk assessments on established and potential invasive species. This has identified a number of high risk species and their implications under the Habitats Directive and Water Framework Directive. However, it was necessary for ISI to prioritise species for management action and as such not all invasive species are currently being addressed in as comprehensive a manner as needed. In summary, the greatest threat will be to freshwater habitats and species which are already under the greatest pressure from invasive species and highly vulnerable to new invasions. Increasing impact of invasive species in aquatic habitats may also impact on the ecological status of waterbodies leading to a failure to meet the objectives of the Water Framework Directive.

There are a number of potential invasive species to Ireland who have not yet become established as they are temperature limited, this particularly applies to fish species such as carp species and Zander. There have been unsuccessful attempts by anglers to introduce a range of non-native fish species into Ireland. It is likely that they could become established in the future after deliberate introduction as summer water temperatures become warm enough for successful breeding. A review of which species are expanding their ranges northwards in continental Europe and Britain would identify new potential invaders to Ireland. However, given the many thousands of species this could include, it will be important to focus on those species which will have a negative impact on biodiversity rather than all species which may extend their range to Ireland in response to a changing climate.

Current management strategies and policies relating to invasive species need to incorporate climate change considerations to a greater extent than they currently do. This is particularly relevant for those sectors that import non-native species such as horticulture, the pet and aquaria trade and aquaculture. However, there is an opportunity to do so in the next phase of Invasive Species Ireland and ensure that climate change is fully integrated into risk assessments, invasive species policy and plans and monitoring programmes. Conversely there is also a need to ensure that conservation policies adopted in response to climate change such as increasing connectivity and assisted migration do not inadvertently facilitate species invasions.

9. Capacity building and extending networks

The Irish Platform for Biodiversity Research is due for imminent renewal (Deirdre Lynn, pers. comm.) so it is timely to consider how such a forum could incorporate climate change and further consider the issues raised in this review. In this section we identify some prospective strategic linkages which would improve information sharing in relation to wider initiatives elsewhere. We also provide a brief summary of the benefits which the suggested network affiliations may be able to provide.

9.1 Biodiversity forum links

In terms of Biodiversity Fora, there are a number of UK and European groups with various platforms for information sharing. One obvious candidate link would be to the UK Biodiversity Research Advisory group (UK BRAG). The stated aims of UK BRAG include:

- Identify, promote and facilitate biodiversity research to support UK and individual country biodiversity action plan commitments;
- Coordinate effective and efficient UK engagement with European biodiversity research issues, fulfilling the role of a national biodiversity research platform;
- Contribute to effective biodiversity research networking in the UK, leading to increased interdisciplinary capacity;
- Support knowledge transfer activities in relation to biodiversity research.
[URL: <http://www.jncc.gov.uk/default.aspx?page=3900>]

There are also links to a wide set of publications which could help inform policy developments in Ireland, as well as wider links to European Networks. One possible candidate European link would be to the European Platform for Biodiversity Research Strategy (EPBRS). The EPBRS is a forum at which natural and social scientists, policy-makers and other stakeholders identify, structure and focus the strategically important research that is essential to:

- Use the components of biodiversity in a sustainable way;
- Maintain ecosystem functions that provide goods and services.
[URL: <http://www.epbrs.org/epbrs/>]

9.2 The case for a biodiversity and climate change research forum and suggested links

Given the recurrent theme running throughout this report and echoed elsewhere of the need for improved data, there is a clear need for a biodiversity and climate change research forum which could be an all-island forum organised under the auspices of the Irish Platform for Biodiversity Research. As well as being an important information sharing platform in its own right, such a forum would facilitate data sharing between the various research groups. There is an obvious and emergent need for a metadata repository to further enable biodiversity impacts research and modelling on scales ranging from the national via the landscape and site to the level of individual species.

These data requirements have been recognised for some time. Thus for example, the European Platform for Biodiversity Research Strategy (EPBRS) recognised a need for;

1. Methods to integrate and aggregate data varying in scale, precision and accuracy to answer local, regional and global research and policy questions.
2. Quality assessments and confidence estimates based on data provenance, characteristics, collection bias and processing.
3. Spatial and temporal sampling strategies to take into account statistical characteristics and cost efficiency.
4. Biodiversity informatics to discover and exploit links between ecological, species, genetic, molecular, economic and social data.
5. Tools and criteria for prioritisation of data collection (EPBRS, 2005).

The formation and maintenance of such a forum would also enable more effective partnership working and funding applications. It is encouraging therefore that the Irish Platform for Biodiversity Research is due for imminent renewal, particularly since this platform feeds into the EPBRS. The re-establishment of such a platform would provide a locus for discussion on e.g. establishing protocols on data sharing, accessibility and standardisation for use across the community in Ireland and in line with the principles established by EPBRS.

Similarly, affiliation to wider biodiversity research networks would help identify emerging themes, initiatives and funding opportunities. For example, there is a cluster of European-funded projects at Agricultural Policy-Induced landscape changes: effects on biodiversity and Ecosystem Services. [URL: <http://agripopes.net/links.htm>]. This includes links to for example, A Long-Term Biodiversity, Ecosystem and Awareness Research Network (ALTER-Net);

- ALTER-Net aims to help deliver on the 2010 target by promoting a better integrated and stronger European biodiversity research capacity resulting in the establishment of a lasting infrastructure for integrated ecosystem research, combining ecological and socio-economic approaches, and with greater emphasis on communication with relevant audiences.
[URL: <http://www.alter-net.info/>]

This in turn provides links to other European research networks such as e.g. the European Long-Term Ecosystem Research Network [URL: <http://www.lter-europe.net/>].

10. Developing an indicator species monitoring network

A powerful information tool in tracking the impacts of climate change is the use of biodiversity indicators. Indicators of biodiversity can be viewed in three categories: structural, compositional and functional (Iremonger *et al.*, 2007). At EU level, the European Commission is developing a headline set of biodiversity indicators with the European Environment Agency, to assess achievement of the 2010 target (EEA, 2007). Further development of indicators is needed at national level to inform the public and decision-makers on biodiversity, the effectiveness of conservation measures and progress made in halting biodiversity loss (EPA, 2008).

10.1 Monitoring networks: terrestrial settings

Although there appear to be numerous possibilities for indicators of the impact of climate change on the Irish environment, a review by Donnely *et al.* (2004) highlighted the difficulties involved in identifying an unambiguous set. Consequently they concluded that:

- the main problems relate to the absence of existing long-term data sets which can be extended into the future and from which trends can be observed;
- in addition, the overwhelming influence of factors other than climate change on particular indicator can reduce their significance and value; and
- currently the most effective impact indicator of climate change in Ireland appears to be phenological observations on tree developmental stages (Donnely *et al.*, 2004).

This has led to further work reviewing the utility of tree phenology as an indicator (Donnely *et al.*, 2006). Some authors have also considered that arctic-alpine associations may be useful a useful indicator generally (Nagy 2003, 2006).

There is consensus that the most climate change vulnerable species in Ireland are likely to be Arctic and Boreal relicts and mountain species (Byrne *et al.*, 2003). While it might be expected that oceanic mountains would be buffered against climatic change by their more limited annual temperature range, by comparison with higher mountains such as the Alps, the nival zone is insufficient in extent to accommodate any potential upward migration of species (Crawford, 2000, 2003). In addition, since many mountain plants are intolerant to competition, fast-growing lowland species with broad altitudinal and ecological ranges are predicted to expand at the cost of slow-growing competition-intolerant species with narrow habitat demands (Korner, 1999, 2003; Klanderud and Birks, 2003).

Consequently, a number of authors have argued (see e.g. Coll *et al.*, 2005 and references therein) that these community associations in maritime uplands are likely to be particularly sensitive to climate change. Therefore, and given the poor assessment reported in Table 3, there is a case to be made for initiating an ongoing monitoring programme of these communities as prospective indicators. However, there is also a need generally for widespread monitoring programmes to be initiated for a host of candidate indicator species and communities.

If we consider that the status of priority species provide an indicator of change for a wide range of ecosystems and hence indirectly biodiversity (EHS, 2008). There is a need to expand monitoring programmes as an early warning system in relation to climate change. Not least since as we have highlighted in Section 4 here, even before we consider climate change and biodiversity impact linkages, it is not possible to provide a comprehensive assessment of all aspects of biodiversity in either the Republic or Northern Ireland due to a lack of baseline monitoring data. These substantial knowledge and data gaps in relation to many of our priority species and habitats compromise our ability to protect them from contemporary pressures and in the absence of confounding issues such as climate change.

In recognition of this issue, the NPWS have in recent years directed considerable resources into baseline monitoring and survey work (Annex 2). Therefore some of the indicators used to assess the conservation status of habitats and species could be used as indicators of climate change; e.g. changes in areas of habitat sensitive to coastal squeeze or changes in species assemblages as a response to climate change (Deirdre Lynn, pers. comm.). Similarly, Bird Watch Ireland and the RSPB have recognised the importance of birds as indicators in the Irish landscape in their ongoing monitoring programmes (e.g. Lynas *et al.*, 2007).

10.2 Monitoring networks: marine settings

The dynamic nature of the marine environment and the ability of species to migrate rapidly in response to changes can make monitoring difficult. Conversely, however, if an effective monitoring network is in place, changes in distribution and to species assemblages can be detected relatively quickly. Similarly, it is recognised that protecting pelagic species that exploit ephemeral hydrographic features and fronts is inherently difficult because these are dynamic features defined by short-lived gradients in water properties (Hyrenbach *et al.*, 2000). It has nonetheless been suggested that prospective Marine Protected Areas (MPAs) could adopt the 'core-buffer' zonation model devised for biosphere reserves (Hyrenbach *et al.*, 2000).

It has also been suggested that an analogous approach to structural and functional connectivity in the terrestrial environment could also be adopted for Ireland's marine environment. Thus for example, when considering the marine environment beyond the littoral region;

- the shelf seas could be viewed as a landscape unit;
- while the shelf edge could be viewed as a corridor through which we can monitor changes in the distribution and abundance ranges of marine organisms, this since there is wide evidence that this area is a migration corridor for many marine organisms at different trophic levels (I. Nardello, pers. comm.).

Monitoring changes in marine species and systems will undoubtedly be difficult and expensive, but it is crucial for our understanding of ecosystem changes in response to climatic drivers. Not least since the marine environment is already subject to a variety of threats, ranging from the loss or degradation of biodiversity and changes in its structure, loss of habitats, contamination by dangerous substances and nutrients and possible future

effects of climate change. These and related pressures have recently been recognized with the implementation of the recent Marine Strategy Framework Directive (EU, 2008). Given that indicators and monitoring are a core component of this directive, we see clear scope for synergies in relation to monitoring the impacts of climate change on marine biodiversity.

11. Concluding vision

Ireland's biodiversity is not a static resource; it is profoundly dynamic as a result of natural processes and events and of human use of the land. The legacy of fragmented landscapes and degraded habitats across our island remind us of the ecological consequences of human history. Any vision of the future in a changed climate must include maintaining and enhancing Ireland's high quality remaining natural environment.

Many of our ongoing environmental problems are at least in part a consequence of decisions confined to particular sectors or restricted areas. While our summary in Section 4 indicates that many of our priority habitats and species are in an already parlous state arising from historical and contemporary pressures, and for many we have no insight into what the impacts of climate change could mean. Targeted research is therefore a fundamental priority. However, and given the poor status reports for many of our priority habitats and species (Section 4), proactive conservation interventions are also urgently required.

Despite the considerable uncertainties surrounding future climate projections (Annex Section 1.1), the rate and complexity of recent change in our climate is beyond our experience and there is consensus that future climates will not be like past climates. The implications of a changing climate for many of our native species are likely to be profound, there is therefore a real need to link and harmonize Ireland's policy response to climate change and biodiversity. In addition, there has to be a review of monitoring to assess whether existing systems are sufficiently sensitive to the effects of a changing climate and identify where new systems may be required.

However, and as we have highlighted here, the extent and complexity of many of the cross-cutting issues requires the involvement of all relevant interests to avoid decisions taken in isolation by a single sector, or with respect to an unduly limited area. Our findings here mirror those of other work. Hence we would re-iterate for example, that despite the significant amount of climate change impact research in relation to biodiversity, there does not appear to be any clear co-ordination of strategic planning within and between the various bodies responsible to address climate change risks and planning for adaptation (Arkell *et al.*, 2007). This also means that the links to other sectors, for example between biodiversity, fisheries, tourism and recreation, are potentially missed (Arkell *et al.*, 2007), although we provisionally address these and some wider communication issues in Section 6 above.

There is a clear need for current biodiversity policy having the capacity to deliver for climate change adaptation and mitigation. However, this is substantially under resourced and, as a result, the limited capacity to deliver for biodiversity will impact negatively on dealing with a changing climate. There is therefore a need to build capacity within competent authorities, both in terms of staff resources and expertise in biodiversity and climate change issues if a coherent cross-sectoral response is to be delivered. Similarly, there has to be a willingness to implement and adapt best practise from elsewhere and apply this on an all-island basis.

There is also a need for education and awareness raising, with a particular focus on the human impact on species and habitats and the scale of the likely impacts of a changing climate (Arkell *et al.*, 2007). Perhaps more importantly, we must engage people with the issues and encourage our young people to think beyond traditional barriers. If we accept that biodiversity is the foundation of ecosystem services to which human well-being is intimately linked (RIA, 2008), then it follows that biodiversity conservation is profoundly linked to wider issues of social and generational equity under projections of climate change. Recent work has emphasized that poverty and the loss of ecosystems and biodiversity are inextricably intertwined, with most of the world's poor heavily reliant on ecosystem services (European Communities, 2008).

If Ireland is to effectively incorporate climate change adaptation and mitigation strategies into National Biodiversity Strategies and Action Plans (Wyse-Jackson, 2008). It follows that there must be improved coherence at national level between various plans and programmes affecting biodiversity, and that decision making at regional and local levels must mirror these high-level commitments for biodiversity (EPA, 2008) (Section 5). While such blithe recommendations are fine in principle and on paper, we have reviewed a number of the confounding practical issues and innate theoretical communication and expectation mismatch difficulties in Section 7.1.

In addition, we have identified an added value in having the various national and local biodiversity platforms forging and maintaining identified strategic links across Europe (Section 9.1). Similarly, we have identified a need for the formation of an Ireland-wide climate change and biodiversity research forum in order to improve information sharing between the appropriate agencies of state and the various research clusters. Again, we have identified wider European research networks to which such a forum could affiliate in order to facilitate engagement in wider European initiatives and to provide access to prospective additional funding streams (Section 9.2).

These sorts of findings do not preclude grass roots engagements and interactions, but rather demand it. Despite the Biodiversity Forum's role in increasing public awareness on biodiversity and a number of other awareness raising campaigns and initiatives across Ireland. A 2007 Euro-barometer report on attitudes of Europeans to biodiversity found that 52 per cent of those surveyed in Ireland had never heard of the term 'biodiversity', while 26 per cent had heard of it but did not know what it meant and only 22 per cent had heard of it and knew what it meant (EPA, 2008).

Similarly, by constructing climate change as a global problem, one that is distanced and un-situated relative to an individual's mental world, we make it easy for citizens to verbalise superficial concern with the problem, but a concern belied by little enthusiasm for behavioural change (Slocum, 2004; Lorenzoni *et al.*, 2007; Hulme, 2008). In doing so we have contributed to conditions that yield psychological dissonance in individuals: the contradictions between what people say about climate change and how they act (Stoll-Kleeman *et al.*, 2001; Hulme, 2008).

If we concede that wilderness is the preservation of the world (Thoreau, 1854) is one starting point on our long journey towards ecological consciousness, and if we consider that climate scenarios are socially contingent products of a post-normal science (*sensu* Hulme and Dessai, 2008) as another view along our continuing journey. By linking climate change and biodiversity issues and actions, and through engaging the citizen in the gulf of debate informing the development of scientific and conservation thinking between these interpretations, we can only better inform all our futures.

Given the already poor state of many of our priority habitats, we do not have time to indulge our need for perfect knowledge, and in many cases we already know what to do. With stewardship and management must come a wider recognition that beauty has a place beyond utility and is neither geographically bounded when we reflect upon nature, nor genetically constrained when we consider biodiversity. At the very least, and if we cannot value intrinsic worth, as our new thinking ponders the importance of ecosystem services in provisioning for the least privileged and poor of this and subsequent generations, we should not add a failure of conscience to our apparent inability to act.

Much has been made recently in both the scientific literature and the media of a 'tipping point' being exceeded in a climate context precipitating rapid climate change. However, in a balanced view, the human tipping points are likely to be more important, with a plethora of development choices and mitigation options available. Therefore while the inertia of the climate system commits us to some component of anthropogenic warming over the next few decades, present human activities have not yet entirely defined the magnitude of later century warming. As a result development pathway choices over the next couple of decades are more likely to influence the possible magnitude of end of century warming. This is not, however, a closing door to complacency, the changes which will impact our biodiversity in the coming decades are already underway. What is required is improved policy planning linked to informed public debate to help better shape our island-wide response to a complex global issue.

12. Recommendations

This review contains a range of recommendations which are summarised here. We consider that integrating biodiversity and climate change adaptation and mitigation policy is an obvious priority and we have identified the development of the National Adaptation Strategy and the vehicle for taking this forward along with some of the integrated cross-sectoral policy needs we have identified here.

Specific recommendations include:

- Improve baseline data on the distribution of species and habitats
- Further develop the evidence base for policy through research combining modelling with experiments
- Carry out a fuller climate change vulnerability assessment for annexed habitats and species
- Identify indicator species, include in the Biodiversity Strategy indicator set and develop a indicator species monitoring network
- Increase the area of land managed for biodiversity and other environmental benefits including areas for buffering and linkage outside the protected area network
- Protect and create habitat features across the land mass to make the island more permeable to biodiversity
- Develop strategic linkages with research fora
- Fully implement the Strategic Environmental Assessment Directive which will strengthen biodiversity and climate considerations in sectoral policies
- Integrate biodiversity and climate change policies in the Republic of Ireland through the National Adaptation Plan
- Harmonise adaptation policies on an all-island basis
- Integrate climate change considerations into management plans for the Natura 2000 network
- Ensure that climate change is fully integrated into invasive species policy and plans and monitoring programmes by including it as a requirement of the next phase of Invasive Species Ireland
- Develop sectoral specific adaptation policies
- Carry out a Strategic Environmental Assessment (SEA) of renewable energy policy
- Evaluate REPS measures for their impacts on biodiversity and effectiveness.
- Sustainability Impact Assessments (SIAs) should be carried out to ensure biodiversity is taken into account in the development of renewable energy resources
- Encourage biofuel crops that do not have a negative impact on biodiversity
- Develop guidelines for regional and local planning that incorporate biodiversity and climate change considerations
- Ensure all existing biodiversity laws, policies and strategies are implemented across the island to create resilient populations of species in healthy habitats.

Acknowledgements

The authors are grateful to Comhar for providing the funding and the platform for undertaking the work. We thank all members of the Biodiversity Forum for providing helpful and constructive input at various stages as the manuscript evolved.

We also thank Deirdre Lynn at NPWS for supplying us with the Figures and Tables from the 2008 Annex report, as well as the monitoring programme overview used in Annex 2 here, and for her helpful suggestions in helping to shape the final manuscript. John Coll would like to thank Peter Wyse-Jackson for his helpful discussion input in shaping the content of parts of the manuscript.


Further thanks are extended to: Caitriona Douglas at NPWS; Karin Dubsy at Coastwatch; Siobhán Egan at Bird Watch, Ireland; Hilary Kirkpatrick at the Ulster Wildlife Trust; James Robinson at RSPB, Northern Ireland; Ilaria Nardello and Ken Whelan at the Marine Institute. All gave freely of their time and expertise in providing suggestions to help shape and improve the content of the final manuscript.

Annex 1: Projected climate changes for Ireland


1.1 Climate model projections

A number of studies have applied selected Intergovernmental Panel on Climate Change (IPCC) Special Report on Emissions Scenarios (SRES) (Nakicenovic *et al.*, 2000) to model climatic changes across Ireland at a regional scale. As viewed by climate researchers, a scenario is a coherent, internally consistent, and plausible description of a possible future state of the world (Carter *et al.*, 1994). A ‘climate scenario’ may form one component of a broader scenario of the future, and yet is itself informed by broader descriptions of the future world, for example, demographic trends, energy prices or greenhouse gas emissions (Hulme and Dessai, 2008).

Despite the different modelling approaches, models and scenario combinations used, there is some concordance of projected changes to mean seasonal temperature for the 2020s across Ireland (Figure 1). However, there are more disparities in the magnitude and sign for some of the projected seasonal precipitation changes for the island (Figure 2). In assessing the possible impacts on biodiversity, it is worth noting that the climate modelling community define the 2020s time-slice as 2021 – 2049. Therefore the median of this period coincides with our aim to focus on possible impacts based on a near term (2030s) horizon scan.



Annex Figure 1: Comparison of mean seasonal temperature change projections (°C relative to 1961-1990) for Ireland by the 2020s from different modelling groups. The spatial variation across the island captured by the different groups has been averaged.



Annex Figure 2: Comparison of mean seasonal precipitation change projections (% change relative to 1961-1990) for Ireland by the 2020s from different modelling groups. The spatial variation across the island captured by the different groups has been averaged.

In part these differences can be attributed to the different methods used, as well as the different scenarios. For example, the Irish Climate Analysis and Research Units (ICARUS) results (Fealy and Sweeney 2003, 2007) are the mean values of A2 and B2 scenario outputs obtained by statistically downscaled outputs from the ensemble mean of three Global Climate Models (GCMs). Whereas the Community Climate Change Consortium for Ireland (C4i) (McGrath *et al.*, 2008) and the United Kingdom Climate Change Impacts Programme 2002 (UKCIP02) (Hulme *et al.*, 2002) outputs are dynamically downscaled outputs obtained from a Regional Climate Model (RCM) nested within a driving GCM.

The difference between the C4i and UKCIP02 results are attributable to the different GCMs and RCMs used. Also, both the C4i and UKCIP02 projections are derived from weighted ensemble averages from the B2 family of scenarios with the emphasis more on local solutions to economic, social and environmental sustainability (Table 1). Whereas the A2 family of scenarios project a greater increase in global carbon dioxide (CO₂) emissions associated with different development trajectory assumptions (Nakicenovic *et al.*, 2000) (Table 1). The ICARUS results hence incorporate some of the greater temperature increases associated with this scenario relative to the more conservative assumptions (in CO₂ emissions terms) of the B2 family of scenarios.

Annex Table 1: Summary linking ICARUS, C4i and UKCIP02 climate change scenarios to other scenario developments (adapted from Hulme *et al.*, 2002)

SRES¹ Storyline	OST² Foresight Scenario	UKCIP98 Socio- economic Scenario	UK Environment Agency Scenario	UKCIP02 Climate change Scenario
B1	Global Sustainability	Global Sustainability	Gamma	Low- Emissions
B2	Local Stewardship	Local Stewardship	Delta	Medium-Low Emissions
A2	Provincial Enterprise	National Enterprise	Alpha	Medium-High Emissions
A1F1	World Markets	World Markets	Beta	High Emissions

Notes:

¹ SRES: Special Report on Emissions Scenarios


² OST: UK Office of Science and Technology

Uncertainty is perhaps the single most sensitive facet of a climate scenario and one that climate science has struggled to come to terms with (Shackley and Wynne, 1996; Van der Sluijs, 1997; Van der Sluijs *et al.*, 1998; Moss and Schneider, 2000; Lahsen, 2005). Arising from this, an essential component of the communication and dissemination process for climate scenarios is to convey the range of assumptions, conflicts and compromises that have been made in their construction (Hulme and Dessai, 2008).

Amid these and other issues, and despite the advances offered by the climate modelling community, there are remaining concerns surrounding the validity of the socio-economic and development trajectory scenario assumptions used to derive the greenhouse gas (GHG) emissions scenarios driving the models. With the IPCC having decided there was no time to develop new scenarios for its 4th assessment report, the various modelling groups are using the same SRES scenarios used in the previous assessment (Schiermeier, 2006). This has led to criticism from many economists on the basis that the existing SRES scenarios rely on outdated economic theories which fail to reflect how lifestyle and energy demand in both rich and poor countries are likely to change (Schiermeier, 2006; Van Vuuren and O'Neill, 2006).

However, these concerns and criticisms largely relate to projections of change for later in the century. There is scientific consensus that the thermal inertia of the climate system commits us to the changes projected for the 2030s based on present GHG levels irrespective of mitigation measures in the intervening decades. Thus, for example, there is close agreement of globally averaged surface air temperature (SAT) multi-model mean warming for the early 21st century for concentrations derived from the three non-mitigated IPCC SRES B1, A1B and A2 scenarios (Meehl *et al.*, 2007). Therefore the warming averaged for 2011 to 2030 is affected little by different scenario assumptions or different model sensitivities (Meehl *et al.*, 2007). As a result, there is considerable

convergence in the projections of climate change from the various models for this decade, with the later century divergence between the model results reflecting the increasing uncertainty (Figure 3).



Annex Figure 3: Multi-model means of surface warming (relative to 1980–1999) for the scenarios A2, A1B and B1, shown as continuations of the 20th-century simulation (Meehl *et al.*, 2007).

Given some of the scenario-related issues outlined here and given that other drivers of change will interact with climate change to impact biodiversity (Section 5), we consider our review here in the context of a near-term (2030s) horizon scan. Also, in recognition that the potential for developing synergies between climate change mitigation and adaptation has become a recent focus for both climate research and policy across sectors (Klein *et al.*, 2005). Here we have attempted to assess how an emergent policy on climate change and biodiversity might facilitate the successful integration and implementation of mitigation and adaptation measures for the island’s biodiversity resource.

1.2 Summary knowledge review: other drivers

1.2.1 Sea level rise

Sea level is determined by changes in water volume or coastal elevation effected by the warming of the water (causing expansion) or by land movement; specifically for Ireland, isostatic readjustment causing land rebound after the last glaciation. Regional differences in climate-induced sea level rise can vary by up to +/- 50% and there is little agreement between climate models about these regional patterns of sea level rise, although Hulme *et al.* (2002) suggest that thermally driven changes in sea-level will be between +0.04 m and +0.14 m by the 2030s. However, while there is agreement that sea levels will rise as the planet warms, the complex dynamic and thermodynamic interactions of ocean, atmosphere and ice sheets are not fully understood.

There is also a considerable thermal inertia between atmospheric and abyssal ocean warming, for example it has been estimated that the modern ocean overturns once every ~1000 years (Adkins and Pasquero, 2004). However, other recent work indicates that dense water formed in both hemispheres is freshening in response to changes in the high latitude freshwater balance and rapidly transmitting the signature of changes in surface climate into the deep ocean (Rintoul, 2007). For a fuller account of the scientific issues and knowledge gaps in an Ireland-specific context, together with the likely impacts on the coastal zone, the interested reader is directed to the review by Fealy (2003).

1.2.2 Thermohaline circulation (THC) stability

Ocean currents cause significant geographical differences in the supply of heat to the atmosphere and regions around the North Atlantic Ocean have a mean annual surface air temperature that is 5–7°C warmer than those at the same latitude in the Pacific (Stocker and Marchal, 2000). This can be attributed to the THC of the Atlantic Ocean moving warm, saline tropical waters northward, the Gulf Stream being part of this basin-scale circulation (Stocker and Marchal, 2000).

These basin-scale changes in the Atlantic Ocean related to the THC have been an important driver of multi-decadal variations in the summertime climate of North America and western Europe (Sutton and Hodson, 2005). This has led a number of authors (Enfield *et al.*, 2001; Mc Cabe *et al.*, 2004; Knight *et al.*, 2005) to suggest that changes in the Atlantic Ocean associated with a pattern of variation known as the Atlantic Multidecadal Oscillation (AMO) were responsible.

The existence of abrupt past climate changes has fuelled concern over the possibility of similar changes in future, particularly if anthropogenic climate change might trigger another instability of the circulation and a severe cooling over the North Atlantic and parts of Europe (Rahmstorf and Ganopolski, 1999; Rahmstorf, 1999, 2000; Srokosz and Gommenginger, 2002; Stouffer *et al.*, 2006). Despite these considerable uncertainties surrounding the stability and behaviour of the THC on varying timescales, consensus remains that a major ocean circulation change should be considered a 'low probability-high impact' risk, but emphasises that proper risk analysis is crucial for this type of non-linear climatic change (Rahmstorf, 2000; Hulme *et al.*, 2002; Schmittner *et al.*, 2005).

While most GCMs simulate a weakening of the THC over the 21st century, none demonstrate a shut down by 2100 and overall warming is predicted to offset any cooling associated with this weakening (Hulme *et al.*, 2002; NERC, 2006).

1.2.3 Changes in storm and wave climate

There is strong evidence for increased wave heights in north east Atlantic waters and for increased occurrence of strong winds over the UK and adjacent regions from the 1960s to the present (Woolf and Coll, 2007). It is unclear whether recent behaviour is driven by climate change or is simply natural variation and whether substantial changes in storminess are likely in the 21st century (Woolf and Coll, 2007). Bacon and Carter (1991) inferred an increase in mean wave height of about 2% per year “over the whole of the North Atlantic in recent years, possibly since 1950” from observational data notably from Seven Stones Light Vessel (1962-1986). Recent analyses of a more extensive data set confirm a significant upward trend in wave heights in the North Atlantic, but only for the last 50 years and embedded within a pattern of multi-decadal variability over more than a century (Gulev and Hasse, 1999; Gulev and Grigorieva, 2004).

Wave heights in the North-East Atlantic and northern North Sea are known (from analysis of *in situ* data, satellite data and model reconstructions) to respond strongly and systematically to the behaviour of the North Atlantic Oscillation (NAO) (Woolf and Coll, 2007), which may also be interpreted as a regional manifestation of a wider Northern Annular Mode (NAM) of climate variability. The recent strong trend in the NAO (towards stormier conditions) is apparently unique in its history, but it is controversial whether this is a response to greenhouse gas forcing (Osborn, 2004). Many GCMs suggest a general trend towards the stormier tendency of NAO/NAM in the 21st century (e.g. Terray *et al.*, 2004; Miller *et al.*, 2006).

Factors related to the wind such as storminess and roughness of the sea are recognised to be very difficult to predict within climate change scenarios, with present confidence in GCM and RCM modelled wind field changes remaining low (Hulme *et al.*, 2002). However, a large number of analyses have now been conducted. Some analyses have focussed on general features of the storm track (strength and position) and related changes in regional indices such as the NAO and NAM (Woolf and Coll, 2007). Wolf and Woolf (2006) have shown the sensitivity of the wave climate to such changes. Other analyses have focussed on the number and intensity of extra-tropical cyclones and it is a feature of the analyses that different approaches suggest significantly different views of future storm climate (Woolf and Coll, 2007).

Many Global Climate Models suggest a general trend towards the stormier tendency of NAO/NAM in the 21st century (e.g. Terray *et al.*, 2004; Kuzmina *et al.*, 2005). Both a stronger storm track and a poleward displacement of that track are common features in studies of this kind (e.g., Miller *et al.*, 2006).

Whereas analyses primarily based on RCMs suggest different and mostly weaker changes in winds and storminess (e.g. Hulme *et al.*, 2002; Hanson *et al.*, 2004; Lozano *et al.*, 2004; Leckebusch *et al.*, 2006). An analysis of outputs from fifteen coupled GCMs forced by enhanced greenhouse warming experiments finds that there is a reduction in the

total number of extra-tropical cyclones but an increase in the number of intense events (Lambert and Fyfe, 2006). With no apparent change in the geographical positions of the storm tracks, they conclude that there is no obvious shift in storm tracks associated with global warming (Lambert and Fyfe, 2006).

Considering these contradictory findings, downscaling via general structural changes in the atmosphere (such as shifts in NAO) may be more suitable for storminess than analysing winds in RCMs. Therefore and given that preference, the shift to stormy conditions suggested by Terray *et al.* (2004) and others should carry more weight than the contrary results from RCMs, but it is debatable (Woolf and Coll, 2006).

Annex 2: An overview of the NPWS monitoring programme

NPWS Research Section

Overview

A program is ongoing to record major activities that impact the integrity of designated sites and the annexed habitats and species which are harboured within the sites. This data is collected by NPWS Conservation Rangers. The EU standardised list of activities is being updated to include climate change impacts.

Many local scale management efforts have follow up monitoring in place to determine the effect of the management measures. Scientific programs to monitor conservation status have also been initiated and are steadily developing for a large proportion of annexed habitats and species listed below. These programs aim to monitor a representative sample of the national resource of the targeted habitats or species, therefore sampling is not restricted to designated areas.

Measures taken to establish monitoring/surveillance

Habitats

The monitoring methods, which have been employed or are being devised, address the attributes required to assess Conservation Status following guidelines agreed at EU level, i.e. Habitat extent, Structure & Functions and Future Prospects (positive and negative impacts). For new baseline monitoring many surveys adapted methods used in the UK to assess condition. Where existing surveys had been completed methods were devised to make full use of the older datasets. Areas are selected for survey to ensure a representative sample of the habitat is surveyed. In some instances methods are being tested on a smaller sub-sample of areas.

Species

For species with a restricted distribution detailed ecological and population studies have been undertaken. For more widespread species presence/absence data is being gathered at point locations across the range of the species. Information from other sectors/agencies that may impact on the conservation status of habitats or species is also collated and used to interpret any changes observed, e.g. forestry, agriculture, water quality.

Habitats & Species Covered by Surveys

Habitats

- Baseline monitoring has been undertaken for 13 dune habitats at c180 locations across Ireland (2005-2006)
- 48 Raised bogs and associated habitats were monitored in 2004/5, these results were compared to a survey carried out in 1995.

- Baseline monitoring is underway for 4 saltmarsh habitats at 130 locations across Ireland (2006-2009)
- Baseline monitoring was undertaken for Calcareous and Nardus grassland in designated sites in 2006.
- The semi-natural grasslands of Offaly, Roscommon, Waterford & Cork were surveyed in 2007-8, this baseline work is being rolled out over the next 5 years.
- The fens of Monaghan were surveyed in 2007 and a national baseline survey is due to be rolled out in 2009.
- The native woodland survey of Ireland collected detailed ecological information of woodlands across Ireland. A monitoring prescription is currently being developed as a result of this survey.
- A detailed research program is underway to determine the best indicators to monitor the conservation status of turloughs.
- Scoping and baseline surveys were initiated in 2008 for Limestone pavement, Juniper scrub, Seacliffs, Floating River Vegetation and Upland habitats
- Baseline surveys have been initiated for tidal mudflats and sandflats and subtidal habitats

Species

There are ongoing national monitoring/surveillance of the following species:

Lesser Horseshoe Bat, Leisler's Bat, Common & Soprano Pipistrelle, Nathusius Bat, Daubenton's bat, Brown Long-eared Bat, Fresh-Water Pearl Mussel, White-Clawed Crayfish, Otter, Salmon, Marsh Saxifrage, Vertigo angustior, Vertigo geyeri, Vertigo moulinsiana, Killarney Fern, Slender Naiad, Petalwort, Shining Sickle Moss & Clubmosses

National baseline monitoring/ surveillance have been completed for:

Irish Hare, Grey Seal, Common Seal, Natterjack toad

Surveillance of species at particular sites/catchments have been carried out for:

River Lamprey, Brook Lamprey, Sea Lamprey, Marsh Fritillary, Cetaceans

Research into the ecology of Myotis bat species has been initiated to determine the best approach to monitoring.

Birds - Recent co-ordinated surveys of Red Grouse, Hen Harrier, Chough and Red-throated Diver have been undertaken. The Countryside Bird Survey and I-WeBS are ongoing.

Data projects

There have been several data gathering programmes including:

- Waterbeetles
- Molluscs

- Spiders
- Bees
- Dragonflies
- Rare plants
- Cetaceans
- Bats
- Vegetation quadrats

Short descriptions for some of the projects are presented on:
<http://www.npws.ie/CurrentResearchProjects/>.

References

- Adkins, J. F., and Pasquero, C. (2004). "Deep Ocean Overturning - Then and Now." *Science* **306** (5699): 1143-1144.
- Adriaensen, F., Chardon, J.P., De Blust, G., Swinnen, E., Villalba, S., Gulinck, H., and Matthysen, E.. (2003). "The application of 'least-cost' modelling as a functional landscape model." *Landscape and Urban Planning* **64**(4): 233-247.
- Albritton, D.L., Meira Filho, L.G., Cubasch, U., Dai, X., Ding, Y., Griggs, D.J., Hewitson, B., Houghton, J.T., Isaksen, I., Karl, T., McFarland, M., Meleshko, V.P., Mitchell, J.F.B., Noguer, M., Nyenzi, B.S., Oppenheimer, M., Penner, J.E., Pollonais, S., Stocker, T. & Trenberth, K.E. (2001). Technical Summary of the Working Group I Report. In Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Inter-governmental Panel on Climate Change, ed. by J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell & C.A. Johnson. Cambridge University Press, Cambridge. pp. 21-83.
- Anderson, K. and Bows, A. (2008). Reframing the climate change challenge in light of post-2000 emission trends. *Philosophical Transactions of the Royal Society A*. **366**, 3863-3882.
- Araujo, M. B., Cabeza, M., Thuiller, W., Hannah, L., Williams, P.H. (2004). "Would climate change drive species out of reserves? An assessment of existing reserve-selection methods." *Global Change Biology* **10**: 1618-1626.
- Arnell, B., Darch, G., and McEntee, P. (Eds) (2007). Preparing for a changing climate in Northern Ireland. Belfast & Edinburgh, Scotland and Northern Ireland Forum for Environmental Research (SNIFFER). Environment and Heritage Service Northern Ireland, Department of Environment Northern Ireland: 282pp.
- Arnott, S. A., and Ruxton, G.D. (2002). "Sandeel recruitment in the North Sea: demographic, climatic and trophic effects." *Marine Ecology Progress Series* **238**: 199-210.
- Bacon, S., and Carter, D.J.T. (1993). "Wave climate changes in the North Atlantic and North Sea." *International Journal of Climatology*, **11**: 545-558.
- Barnett, T., Zwiers, F., Hegerl, G., Allen, M., Crowley, T., Gillett, N., Hasselmann, K., Jones, P., Santer, B., Shnur, R., Stott, P., Taylor, P., Tett, S. (2005). "Detecting and Attributing External Influences on the Climate System: A Review of Recent Advances." *Journal of Climate* **18**: 1291-1314.
- Bennet, S., Sheate, W.R. (2007). The Water Framework Directive, Assessment, Participation and Protected Areas: What are the Relationships? Wexford, Environmental Protection Agency: pp118.
- Berry, P. M., Dawson, T.P., Harrison, P.A., and Pearson, R.G. (2002). "Modelling potential impacts of climate change on the bioclimatic envelope of species in Britain and Ireland." *Global Ecology and Biogeography* **11**: 453-462.
- Berry, P. M., Harrison, P.A., Dawson, T.P. and Walmsley, C.A. (Eds) (2005). Modelling Natural Resource Responses to Climate Change (MONARCH). A Local Approach:

- Development of a conceptual and methodological framework for universal application. Oxford: 277pp.
- Birks, H. J. B. (1997). Scottish biodiversity in a historical context. Biodiversity in Scotland: Status, Trends and Initiatives. L. V. Fleming, Newton, A.C., Vickery, J.A. & Usher, M.B. (Eds.). Edinburgh, The Stationery Office: 21-35.
- Bolens, R., Monchin, D., and O'Sullivan, G. (2005). Climate Change - Implications for Ireland's Marine Environment and Resources: Marine Foresight Series No. 2. Galway, Marine Institute: pp40.
- Bomhard, B., Richardson, D.M., Donaldson, J.S., Hughes, G.O., Midgley, G.F., Raimondo, D.C., Rebelo, A.G., Rouget, M., Thuiller, W. (2005). "Potential impacts of future land use and climate change on the red list status of the Proteaceae in the Cape Floristic region, South Africa." *Global Change Biology* **11**: 1452-1468.
- Buckley, P. (2004). "Development Of Wildlife Conservation Policy And Legislation In Ireland." *Biology And Environment: Proceedings Of The Royal Irish Academy* **104B** (3): 107-111.
- Bullock, C., Kretsch, C., Candon, E. (2008). The Economic and Social Aspects of Biodiversity: Benefits and Costs of Biodiversity in Ireland. Dublin, Environment, Heritage and local Government: pp196.
- Burns, C. E., Johnson, K.M., Schmitz, O.J. (2003). "Global climate change and mammalian species diversity in US national parks." *Proceedings of the National Academy of Science USA* **100**: 11474-11477.
- Byrne, C., Jones, M., and Donnelly, A., Wilson, J. (2003). Assessment of the Impacts of Climate Change on Biodiversity in Ireland. Climate Change Scenarios and Impacts For Ireland. J. Sweeney, (Ed.). Dublin, Environmental Protection Agency. ERTRI Report 15: 247pp.
- Cabeza, M., Moilanen, A. (2001). "Design of reserve networks and the persistence of biodiversity." *Trends in Ecology and Evolution* **16**: 242-248.
- CAP (2006). Ireland CAP Rural Development Programme 2007-2013. Dublin, CAP Rural Development Division: pp202.
- Capdevila –Arguelles, L. and Zilletti, B. (2008). A perspective on climate change and invasive alien species. Council of Europe. T-PVS/Inf (2008) 5.
- Carter, T. R., Parry, M.L., Harasawa, H., Nishioka, S. (1994). IPCC Technical Guidelines for Assessing Climate Change Impacts and Adaptations. London/Tsubuka, UCL/CGER: pp59.
- Carter, T. R., Hulme, M. and Viner, D. (Eds.) (1999). Representing Uncertainty in Climate Change Scenarios and Impact Studies. ECLAT-2, Report No.1, Helsinki Workshop, 14-16 April 1999, CRU, Norwich, UK.: pp128.
- CFB (2000). Central and Regional Fisheries Boards: Annual Report 1999. Dublin, Central Fisheries Board: pp32.
- Charlton, R. A., and Moore, S. (2003). The Impact of Climate Change on Water Resources in Ireland. Climate Change Scenarios and Impacts For Ireland. J. Sweeney, (Ed.). Dublin, Environmental Protection Agency. ERTRI Report 15: 247pp.

- Charlton, R., Fealy, R., Moore, S., Sweeney, J., and Murphy, C. (2006). "Assessing the impact of climate change on water supply and flood hazard in Ireland using statistical downscaling and hydrological modelling techniques." *Climatic Change* **74**: 475-491
- Clerkin, S. (2002). "Incorporating Biodiversity Considerations In Policy." *Biology and Environment: Proceedings of the Royal Irish Academy* **102B (3)**: 177-182.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., De Noblet, N., Friend, A. D., Friedlingstein, P., Grinwald, T., Heinesch, B., Keronen, P., Knohl, A., Krinner, G., Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J. M., Papale, D., Pilegaard, K., Rambal, S., Seufert, G., Soussana, J. F., Sanz, M. J., Schulze, E. D., Vesala, T., Valentini, R. (2005). "An unprecedented reduction in the primary productivity of Europe during 2003 caused by heat and drought." *Nature* **437**: 529-537.
- Clenaghan, C. (2008). Chapter 13: Biodiversity. Ireland's Environment 2008. EPA. Wexford, Environmental Protection Agency: pp273.
- Clerkin, S. (2002). "Incorporating Biodiversity Considerations In Policy." *Biology and Environment: Proceedings of the Royal Irish Academy* **102B(3)**: 177-182.
- Coll, J., Gibb, S.W., and Harrison, J. (2005). Modelling future climates in the Scottish Highlands - an approach integrating local climatic variables and regional climate model outputs. Mountains Of Northern Europe: Conservation, Management, People and Nature. D. B. A. Thompson, Price, M.F., and Galbraith, C.A. (Eds). Edinburgh, TSO, Scotland: 103-119.
- Council of the Parties: UNEP (2008). COP 9 Decision IX/2: pp5.
- Craig, A. (2001). "The Role of The State In Protecting Natural Areas In Ireland: 30 Years Of Progress." *Biology and Environment: Proceedings of the Royal Irish Academy* **101B (1-2)**: 141-149.
- Crooks, K. R., and M., Sanjayan. (2006). "Connectivity conservation: maintaining connections for nature." Connectivity Conservation. Crooks, R. and, M., Sanjayan (Eds.). Cambridge, Cambridge University Press: 1-19.
- Cross, J. R. (2006). "The Potential Natural Vegetation Of Ireland." *Biology and Environment: Proceedings of the Royal Irish Academy* **106B (2)**: 65-116.
- Crawford, R. M. M. (2000). "Ecological hazards of oceanic environments (Tansley Review No. 114)." *New Phytologist* **147**: 257-281.
- Crawford, R. M. M. (2000). "Plant Community Responses to Scotland's Changing Environment." *Botanical Journal of Scotland* **53(2)**: 77-105.
- Crawford, R. M. M., Jeffree, C.E. and Rees, W.G. (2003). "Paludification and Forest Retreat in Northern Oceanic Environments." *Annals of Botany* **91**: 213-226.
- Cubasch, U., Meehl, G A, Boer, G J, Stouffer, R J, Dix, N, Noda, A, Senior, C A, Raper, S. & Yap, K S (2001). Projections of future climate change. Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Inter-governmental Panel on Climate Change. J. T. Houghton, Ding, Y., Griggs D.J., Noguer M., van der Linden P.J., Dai, X., Maskell K. and Johnson C.A. (Eds). Cambridge, Cambridge University Press: 525-582.
- DOEHLG (2007). Ireland National Climate Change Strategy 2007-2012. Department of Environment, Heritage and Local Government. Dublin: pp61.

- Donald, P. F., and Evans, A.D. (2006). "Habitat connectivity and matrix restoration; the wider implications of agri-environment schemes." *Journal of Applied Ecology* **43**(2): 209-218.
- Donnelly, A., Jones, M.B., Sweeney, J. (2004). "A review of indicators of climate change for use in Ireland." *International Journal of Biometeorology* **49**: 1-12.
- Donnelly, A., Caffarra, A., Albanito, F., and Jones, M. (2007). The Impact of Climate Change on Semi-Natural Ecosystems in Ireland (Chapter 5). In: *Climate Change in Ireland: Refining the Impacts*. Dublin, Environmental Protection Agency: pp192.
- Drinkwater, K. F., Belgrano, A., Borja, A., Conversi, A., Edwards, M., Greene, C.H., Ottersen, E., Pershing, A. J., and Walker, H. (2003). The Response of Marine Ecosystems to Climate Variability Associated With The North Atlantic Oscillation. The North Atlantic Oscillation, Climatic Significance and Environmental Impact. J. W. Hurrell, Kushnir, Y., Ottersen, G., Visbeck, M. Washington DC, American Geophysical Union. **134**: 211-234.
- Dunne, S., Hanafin, J., Lynch, P., McGrath, R., Nishimura, E., Nolan, P., Ratnam, J.V., Semmler, T., Sweeney, C. and Wang, S. (2008). Ireland in a Warmer World. Scientific Predictions of the Irish Climate in the Twenty-First Century. R. McGrath, Lynch, P. (Eds.). Dublin, Community Climate Change Consortium for Ireland (C4I): pp109.
- EEA (2007). *Halting the Loss of Biodiversity by 2010: Proposal for a First Set of Indicators to Monitor Progress in Europe*. Copenhagen, European Environment Agency.
- EEA (2008). *Impacts of Europe's changing climate - 2008 indicator-based assessment*. Copenhagen, Joint EEA-JRC-WHO report: pp247.
- EHS (2008). 'Our Environment, Our Heritage, Our Future'. *State of the Environment Report for Northern Ireland*. Belfast, Environment and Heritage Service: pp70.
- Emblow, C. Minchin, D. Mayes, E. Wilson, J. (2003). *Climate Change and the Irish Marine Environment. Climate Change Scenarios and Impacts For Ireland*. J. Sweeney, (Ed.). Dublin, Environmental Protection Agency. ERTRI Report 15: 247pp.
- Enfield, D. B., Mestaz-Nunez, A.M., Trimble, P.J. (2001). "The Atlantic Multidecadal Oscillation and its relationship to rainfall and river flows in the continental U.S." *Geophysical Research Letters* **28**: 2077-2080.
- EPA (2008). Ireland's Environment 2008. Wexford, Environmental Protection Agency: pp273.
- EPBRS (2005). *Draft Recommendations Of The Workshop Of The European Platform For Biodiversity Research Strategy. Innovative Research For Biodiversity Data Acquisition & Assessment*, Brussels.
- Erhard, M., van Minnen, J., Voight, T. (2002). Proposed core set of climate change state and impact indicators in Europe. Copenhagen, European Environment Agency.
- European Commission (2007) *Adapting to climate change in Europe – options for EU action*. SEC(2007) 849.
- European Communities (2008). The Economics of Ecosystems and Biodiversity. An Interim Report. Brussels, European Communities: pp65
- EU (2008). *Marine Strategy Framework Directive*. 2008/56/EC: pp22.

- Evans, C. D., Monteith, D.T., Cooper, D.M. (2005). "Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts." *Environmental Pollution* **137**: 55-71.
- Fealy, R., and Sweeney, J. (2003). Climate Scenarios for Ireland. Climate Change Scenarios and Impacts for Ireland. J. Sweeney. Dublin, Environmental Protection Agency: pp247.
- Fealy, R. (2003). The Impacts of Climate Change on Sea Level and the Irish Coast. Climate Change Scenarios and Impacts For Ireland. In Climate Change Scenarios and Impacts for Ireland. J. Sweeney, (Ed.). Dublin, Environmental Protection Agency. ERTRI Report 15: 247pp.
- Fealy, R., and Sweeney, J. (2007). "Statistical downscaling of precipitation for a selection of sites in Ireland employing a generalised linear modelling approach." *International Journal of Climatology* **27**: 2083-2094.
- Ferris, R. E. (2007). Research Needs for UK Biodiversity. A summary of the important knowledge gaps, identified by the UK Biodiversity Research Advisory Group, 2003-2006. London, Department for Environment, Food and Rural Affairs: pp44.
- Foden, W., Midgley, G.F., Hughes, G.O., Bond, W.J., Thuiller, W., Hoffman, N.T., Kaleme, P., Underhill, L.G., Rebelo, A.G., Hannaah, L. (2007). "A changing climate is eroding the geographical range of the Namib Desert tree Aloe through population declines and dispersal lags." *Diversity and Distribution* **13**: 645-653.
- Frederiksen, M., Harris, M.P., Daunt, F., Rothery, P., and Wanless, S. (2004a). "Scale-dependant climate signals drive breeding phenology of three seabird species." *Global Change Biology* **10**: 1214-1221.
- Frederiksen, M., Wanless, S., Harris, M.P., Rothery, P. and Wanless, S. (2004b). "The role of industrial fishery and oceanographic changes in the decline of the North Sea black-legged kittiwakes." *Journal of Applied Ecology* **41**(6): 1129-1139.
- FSBI (2007) Climate change and the fishes of Britain and Ireland: Briefing Paper 4. Fisheries Society of the British Isles, Granta Information Systems, 82A High Street, Sawston, Cambridge, CB2 4H, UK.
- Gardiner, S., Hanson, S., Nicholls, R., Zhang, Z., Jude, S., Jones, A., Richards, J., Williams, A., Spencer, T., Cope, S., Gorczynska, M., Bradbury, A., McInnes, R., Ingleby, A., and Dalton, H. (2007). "The Habitats Directive, Coastal Habitats and Climate Change - Case Studies from the South Coast of the UK." Southampton, Tyndall Centre for Climate Change Research: pp17.
- Giorgi, F. (2005). "Climate Change prediction." *Climatic Change* **73**: 239-265.
- Giorgi, F. (2006). "Climate change hot-spots." *Geophysical Research Letters* **33** (L08707): doi: 10.1029/2006GL025734.
- Gorham, E. (1991). "Northern peatlands: role in the carbon cycle and probable responses to climatic warming." *Ecological Applications* **1**: 182-195.
- Grosbois, V., and Thompson, P.M. (2005). "North Atlantic climate variation influences survival in adult fulmars." *Oikos* **109**: 273-290.
- Gulev, S. K., and Grigorieva, V. (1999). "Changes of wind waves in the North Atlantic over the last 30 years." *International Journal of Climatology*, **19**: 1091-1117.

- Gulev, S. K., and Grigorieva, V. (2004). "Last century changes in ocean wind wave height from global visual wave data." *Geophysical Research Letters*, **31**: L24302, doi:10.1029/2004GL021040.
- Hannah, L., G. Midgley, et al. (2007). "Protected area needs in a changing climate." *Frontiers in Ecology and the Environment* **5 (3)**: 131-138.
- Hanson, C. E., Holt, T., and Palutikof, J.P. (2004). An Integrated Assessment of the Potential for Change in Storm Activity over Europe: Implications for Insurance and Forestry in the UK. Norwich, Tyndall Centre for Climate Change Research, Final Technical Report IT1.4.; 101pp.
- Harris, M. P. *et al.* (2005). "Effect of wintering area and climate on the survival of adult Atlantic puffins *Fratercula arctica* in the eastern Atlantic." *Marine Ecology Progress Series* **297**: 283-296.
- Harrison, P. A., Berry, P.M. and Dawson, T.P. (Eds.) (2001). Climate Change and Nature Conservation in Britain and Ireland: Modelling natural resource responses to climate change (the MONARCH project). Oxford, UKCIP Technical Report: 1-271.
- Harrison, P. A., Berry, P.M., Henriques, C., Holman, I.P. (2008). "Impacts of socio-economic and climate change scenarios on wetlands: linking water resource and biodiversity meta-models." *Climatic Change* **90**: 113-119.
- Hartig, E. K., Grozev, O., Rosenzweig, C. (1997). "Climate change, agriculture and wetlands in Eastern Europe: Vulnerability, adaptation and policy." *Climatic Change* **36(1-2)**: 107-121.
- Hastie, L.C., Boon, P.J., Young, M.R. & Way, S. (2001). "The effects of a major flood on an endangered freshwater mussel population." *Biological Conservation* **98**, 107-115.
- Hastie, L. C., Cosgrove, P.J., Ellis, N., and Gaywood, M.J. (2003). "The Threat of Climate Change to Freshwater Pearl Mussel Populations." *Ambio* **32 (1)**: 40-46.
- Hegerl, G. C., Stott, P.A., Allen, M.R., Mitchell, J.F.B., Tett, S.F.B., Cubasch, U. (2000). "Optimal detection and attribution of climate change: sensitivity of results to climate model differences." *Climate Dynamics* **16**: 737-754.
- Hill, J. K., Thomas, C.D., and Huntley, B. (2003). "Modelling present and potential future ranges of European butterflies using climate response surfaces." Butterflies, Ecology and evolution taking flight. C. Bogs, Watt, W., and Ehrlich, P. (Eds.). Chicago, Illinois, The University of Chicago Press: 149-167.
- Hilty, J. A., Lidicker, W.J., *et al.* (2006). Corridor ecology: The science and practice of linking landscapes for biodiversity conservation. Washington, Island Press.
- Hopkins, J. J., Allison, H.M., Walmsley, C.A., Gaywood, M. & Thurgate, G. (2007). Conserving biodiversity in a changing climate: guidance on building capacity to adapt. Department for the Environment, Food and Rural Affairs.
- Hoyt, E. (2000). Whale Watching 2000, Worldwide tourism numbers, expenditures and expanding socioeconomic benefits. Bath, UK, Whale and Dolphin Conservation Society: pp36.
- Hulme, M., Jenkins, G.J., Lu, X., Turnpenny, J.R., Mitchell, T.D., Jones, R.G., Lowe, J., Murphy, J.M., Hassell, D., Boorman, P., McDonald, R. and Hill, S. (2002). Climate Change Scenarios for the United Kingdom: The UKCIP02 Scientific Report.

- Norwich, UK., Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia.: 120pp.
- Hulme, M., Dessai, S. (2008). "Negotiating future climates for public policy: a critical assessment of the development of climate scenarios for the UK." *Environmental Science and Policy* **11**: 54-70.
- Hulme, M. (2008). "Geographical Work at the Boundaries of Climate Change." *Transactions of the Institute of British Geographers* **33**(1): 5-11.
- Huntley, B., Green, R.E., Collingham, Y.C., Willis, S.G. (2007). A Climatic Atlas of European Breeding Birds. Cambridge, Durham University, the RSPB, Lynx Edicions.
- Hyrenbach, D. K., Forney, K.A., and Dayton, P.K. (2000). "Marine protected areas and ocean basin management (Viewpoint)." *Aquatic Conservation: Marine and Freshwater Ecosystems* **10**: 437-458.
- IPCC, (2007): Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp. 996
- Iremonger, S., O'Halloran, J., Kelly, D.L., Wilson, M.W., Smith, G.F., Gittings, T., Giller, P.S., Mitchell, F.J.G., Oxbrough, A., Coote, I., French, L., O'Donoghue, S., McKee, A., M., Pithon, J., and A. O'Sullivan, Neville, P., O'Donnell, V., Cummins, V., Kelly, T.C., and Dowding, P. (2007). Biodiversity in Irish Plantation Forests. Wexford, Environmental Protection Agency: pp52.
- Jones, M. B., Donnelly, A., and Albanito, F. (2006). "Responses of Irish vegetation to future climate change." *Biology and Environment: Proceedings of the Royal Irish Academy* **106B** (3): 323-334.
- Kerr, A., and Ellis, N., (2001) 'Managing the impacts of climate change on the natural environment.' In Earth Science and The Natural Heritage: Interactions and Integrated Management., J.E. Gordon, and Leys, K.F., (Eds.). Stationery Office Edinburgh.: Edinburgh. p. 239-250.
- Kettunen, M, Terry, A., Tucker, G. & Jones A. (2007) Guidance on the maintenance of landscape features of major importance for wild flora and fauna - Guidance on the implementation of Article 3 of the Birds Directive (79/409/EEC) and Article 10 of the Habitats Directive (92/43/EEC). Institute for European Environmental Policy (IEEP).
- Kirkpatrick, A. H. (2008). "Building a better future for the uplands in Northern Ireland." *Aspects of Applied Biology* **85**(Shaping a vision for the Uplands): 1-6.
- Klanderud, K., and Birks, H. J. B. (2003). "Recent increases in species richness and shifts in altitudinal distributions of Norwegian mountain plants." *Holocene* **13**: 1-6.
- Klein, R. T., Schipper, E.L.F., Dessai, S. (2005). "Integrating mitigation and adaptation into climate and development policy: three research questions." *Environmental Science and Policy* **8**: 579-588.Brussels, 114 pp. & Annexes.
- Knight, J. R., Allan, R.J., Folland, C.K., Vellinga, M., Mann, M.E. (2005). "A signature of persistent natural thermohaline circulation cycles in observed climate." *Geophysical Research Letters* **32**: L20708. doi:10.1029/2005GL024233.
- Korner, C. (1999). Alpine plant life. Functional plant ecology of high mountain ecosystems. Berlin, Springer.

- Korner, C. H., Paulsen, J., and Pelaez-Riedl, S. (2003). "A Bioclimatic Characterisation of Europe's Alpine Areas." *Alpine Biodiversity in Europe*. L. Nagy, Grabherr, G., Korner, C., Thompson, D.B.A. (Eds.). Berlin, Springer-Verlag: pp13-28.
- Kuzmina, S. I., Bengtsson, L., Johannessen, O.M., Drange, H., Bobylev, L.P., and Miles, M.W. (2005). "The North Atlantic Oscillation and greenhouse gas forcing." *Geophysical Research Letters* **32**: L04073.
- Laffoley, D. *et al.* (2005). *The MarClim Project. Key messages for decision makers and policy advisers and recommendations for future administrative arrangements and management measures*. English Nature Research report, No. 671.
- Lahsen, M. (2005). "Seductive simulations? Uncertainty distribution around climate models." *Social Studies of Science* **35/6**: 895-922.
- Lambert, S. J., Fyfe, J.C. (2006). "Changes in winter cyclone frequencies and strengths simulated in enhanced greenhouse warming experiments: results from the models participating in the IPCC diagnostic exercise." *Climate Dynamics*. **26**: 713-728.
- Lavergne, S., Thuiller, W., Molina, J., Debussche, M. (2005). "Environmental and human factors influencing rare plant local occurrence, extinctions and persistence: a 115 year study in the Mediterranean region." *Journal of Biogeography* **32**: 799-811.
- Leckebusch, G. C., Koffi, B., Ulbrich, U., Pinto, J.G., Spanghel, T., Zacharias, S. (2006). "Analysis of frequency and intensity of European winter storm events from a multi-model perspective, at synoptic and regional scales." *Climate Research*, **31**: 59-74.
- Lee, M. (2001). "Coastal defence and the habitats directive: predictions of habitat change in England and Wales." *The Geographical Journal* **167**(1): 39-56.
- Lorenzoni, I., Nicholson-Cole, S., and Whitmarsh, L. (2007). "Barriers perceived to engaging with climate change among the UK public and their policy implications." *Global Environmental Change* **17**(3/4): 445-459
- Lozano, I., Devoy, R.J.N., May, W., Andersen, U. (2004). "Storminess and vulnerability along the Atlantic coastlines of Europe: analysis of storm records and of a greenhouse gases induced climate scenario." *Marine Geology* **210**: 205-225.
- Lynas, P., Newton, S.F., and Robinson, J.A. (2007). "The status of birds in Ireland: an analysis of conservation concern 2008-2013." *Irish Birds* **8**: 149-167.
- McAree, D. (2002). "The Forest Service Biodiversity Plan." *Biology And Environment: Proceedings Of The Royal Irish Academy* **102B**: 183-184.
- McCabe, G. J., Palecki, M.A., Betancourt, J.L. (2004). "Pacific and Atlantic Ocean influences on multidecadal drought frequency in the United States." *Proceedings of the National Academy of Science* **101**: 4136-4141.
- MCCIP (2008). *Newsletter, Issue 11, August 2008*. Lowestoft, Marine Climate Change Impacts Partnership: pp5.
- McElwain, L., and Sweeney, J. (2007). *Key meteorological indicators for climate change in Ireland*. Wexford, Environmental Protection Agency: pp31.
- McMahon, B. J., Purvis, G., and Whelan, J. (2008). "The influence of habitat heterogeneity on bird diversity in Irish farmland." *Biology and Environment: Proceedings of the Royal Irish Academy* **108B**(1): 1-8.

- Meehl, G. A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, and A. J. W. a. Z.-C. Z. I.G. Watterson (2007). Global Climate Projections. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). Cambridge, Cambridge University Press.
- Meinshausen, M. 2006 What does a 2°C target mean for greenhouse gas concentrations? A brief analysis based on multi-gas emission pathways and several climate sensitivity uncertainty estimates. In *Avoiding dangerous climate change* (eds H. J. Schellnhuber, W. Cramer, N. Nakicenovic, T. Wigley & G. Yohe), pp. 253–279. Cambridge, UK: Cambridge University Press.
- Menzel, A., Fabian, P. (1999). "Growing season extended in Europe." *Nature* **397**: 659.
- Millennium Ecosystem Assessment. (2005). Ecosystems and human well-being: synthesis. Washington, DC, Island.
- Miller, R. L., Schmidt, G.A. and Shindell, D.T. (2006). "Forced annular variations in the 20th century Intergovernmental Panel on Climate Change Fourth Assessment Report models." *Journal of Geophysical Research* **111**: D18101, doi:10.1029/2005JD006323.
- Mitchell, R. J., Morecroft, M.D., Acreman, M., Crick, H.Q.P., Frost, M., M.Harley, Maclean, I.M.D., Mountford, O., Piper, J., Pontier, H., Rehfisch, M.M., Ross, L.C., Smithers, R.J., Stott, A., Walmsley, C.A., Watts, O., and Wilson, E. (2007). England Biodiversity Strategy - Towards adaptation to climate change. Final Report to Defra for contract CR0327. London, Department for the Environment, Food and Rural Affairs.
- Mitsch, W. J., Gosselink, J.G. (1993). Wetlands. New York, Van Nostrand Reinhold.
- Moss, R., Schneider, S. (2000). Uncertainties pp 33-52 in, Cross-cutting guidance papers: IPCC supporting materials. R. Pachauri, Tanguchi, T., Tanaka, K. (Eds.). Geneva.
- Murphy, H. T., and Lovett-Doust, J. (2004). "Context and connectivity in plant metapopulations and landscape mosaics: does the matrix matter?" *Oikos* **105**: 3-14.
- Nagy, L., Grabherr, G., Korner, C., Thompson, D.B.A. (2003). "Alpine Biodiversity in Space and Time: a Synthesis." Alpine Biodiversity in Europe. L. Nagy, Grabherr, G., Korner, C., Thompson, D.B.A. (Eds.). Berlin, Springer-Verlag: 453-464.
- Nagy, L. (2003). The High Mountain Vegetation of Scotland. Alpine Biodiversity in Europe. L. Nagy, Grabherr, G., Korner, C., Thompson, D.B.A. (Eds.). Berlin, Springer-Verlag: pp39-46.
- Nagy, L. (2006). "European High Mountain (Alpine) Vegetation and Its Suitability for Indicating Climate Change Impacts." *Biology and Environment: Proceedings of the Royal Irish Academy* **106B**(3): 335-341.
- Nakicenovic, N., Alcamo, J., Davis, G., de Vries, B., Fenhann, J., Gaffin, S., Gregory, K., Grubler, A., Yong Jung, T., Kram, T., Lebre La Rovere, E., Michaelis, L., Mori, S., Morita, T., Pepper, W., Pitcher, H., Price, L., Riahi, K., Roehrl, A., Rogner, H.H., Sankovski, A., Schlesinger, M., Shukla, P., Smith, S., Swart, R., van Rooijen, S., Victor, N., Dadi, Z. (2000). Summary for Policy Makers: Emissions Scenarios. A

Special Report of IPCC Working Group III. Intergovernmental Panel on Climate Change: pp27.

- NERC (2006). Rapid Climate Change Science Plan. Southampton.
<http://www.noc.soton.ac.uk/rapid/sci/Scienceplan.php>, Natural Environment Research Council: pp15.
- Nicholls, R. J., and Wilson, T. (2001). Integrated impacts on coastal areas and river flooding. Regional Climate Change Impact and Response Studies in East Anglia and North West England(RegIS). I. P. a. L. Holman, P. J. (Eds.). Oxford, UK Climate Impacts Programme (UKCIP): 54-103.
- NPBR (2003). Biodiversity Knowledge Programme for Ireland. Wexford, National Platform for Biodiversity Research: pp30.
- NPWS (2008). The Status of EU Protected Habitats and Species in Ireland. Dublin, National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government: pp135.
- Osborn, T. J. (2004). "Simulating the winter North Atlantic Oscillation: the roles of internal variability and greenhouse gas forcing." *Climate Dynamics* **22**: 605-623.
- Parnesan, C. (2007). "Influences of species, latitudes and methodologies on estimates of phenological response to global warming." *Global Change Biology* **13**: 1860-1872.
- Parnesan, C., Yohe, G. (2003). "A globally coherent fingerprint of climate change impacts across natural systems." *Nature* **421**: 37-42.
- Pethick, J. (2002). "Estuarine and tidal wetland restoration in the United Kingdom: Policy versus practice." *Restoration Ecology* **10** (3): 431-437.
- Pounds, A. J., Bustamante, M.R., Coloma, L.A., Consuegra, J.A., Fogden, M.P.L., Foster, P.N., La Marca, E., Masters, K.L., Merino-Viteri, A., Puschendorf, R., Ron, S.R., Sanchez-Azofeifa, G.A., Still, C.J., Young, B.E. (2006). "Widespread amphibian extinctions from epidemic disease driven by global warming." *Nature* **439**: 161-167.
- Pressey, R. L., Cabeza, M., Watts, M.E., Cowling, R.M., Wilson, K.A. (2007). "Conservation planning in a changing world." *Trends in Ecology and Evolution* **22** (11): 583-592.
- Rahmstorf, S. (1999). "Shifting seas in the greenhouse?" *Nature* **399** (6736): 523-524.
- Rahmstorf, S., and Ganopolski, A. (1999). "Simple theoretical model may explain apparent climate instability." *Journal of Climate* **12** (5): 1349-1352.
- Rahmstorf, S. (2000). "The thermohaline ocean circulation: A system with dangerous thresholds? An editorial comment." *Climatic Change* **46** (3): 247-256.
- RIA (2008). 6th Scientific Statement: Impacts of Climate Change on Biodiversity. Dublin, Royal Irish Academy: pp2.
- Richards, J. A., Mokrech, M., Berry, P.M., Nicholls, R.J. (2008). "Regional assessment of climate change impacts on coastal and fluvial ecosystems and the scope for adaptation." *Climatic Change* **90**: 141-167.
- Rindorf, A., Wanless, S., and Harris, M.P. (2000). "Effects of changes in sandeel availability on the reproductive output of seabirds." *Marine Ecology Progress Series* **202**: 241-252.

- Rintoul, S. R. (2007). "Rapid freshening of Antarctic Bottom Water formed in the Indian and Pacific oceans." *Geophysical Research Letters* **34** (6): L06606.
- Root, T. L., MacMynowski, D.P., Mastrandrea, M.D., Schneider, S.H. (2005). "Human-modified temperatures induce species changes: joint attribution." *Proceedings of the National Academy of Science USA* **102**: 7465-7469.
- Schiermeier, Q. (2006). "The costs of global warming." *Nature* **439**: 375-377.
- Schmittner, A., Latif, M., Schneider, B. (2005). "Model projections of the North Atlantic thermohaline circulation for the 21st century assessed by observations." *Geophysical Research Letters* **32** (23): Art. No. L23710.
- Shackley, S., Wynne, B. (1996). "Representing uncertainty in global climate global climate change science and policy: boundary ordering devices and authority." *Science, Technology and Human Values* **21**: 275-302.
- Slocum, R. (2004). "Polar bears and energy-efficient lightbulbs: strategies to bring climate change home." *Environment & Planning D: Society and Space* **22**: 413-438.
- Srokosz, J., Gommenginger, C. (2002). NERC 'RAPID' Programme Science Plan. Southampton, Natural Environmental Research Council: 15pp.
- Stocker, T. F., and Marchal, O. (2000). "Abrupt climate change in the computer: is it real?" *Proceedings of the National Academy of Sciences of the United States of America* **97** (4): 1362-1365.
- Stoll-Kleeman, S., O'Riordan, T., and Jaeger, C.C. (2001). "The psychology of denial concerning climate mitigation measures: evidence from Swiss focus groups." *Global Environmental Change* **11**: 107-117.
- Stouffer, R. J., Yin, J., Gregory, J.M., Dixon, K.W., Spelman, M.J., Hurlin, W., Weaver, A.J., Eby, M., Flato, G.M., Hasumi, H., Hu, A., Jungclaus, J.H., Kamenkovich, I.V., Levermann, A., Montoya, M., Murakami, S., Nawrath, S., Oka, A., Peltier, W.R., (2006). "Investigating the causes of the response of the thermohaline circulation to past and future climate changes." *Journal of Climate* **19** (8): 1365-1387.
- Sutherland, W. J., Armstrong-Brown, S., Armsworth, P.R., Breerton, T., Brickland, J., C. D. Campbell, Chamberlain, D.E., Cooke, A.I., Dulvy, N.K., Dusic, N.R., Fitton, et al. (2006). "The identification of 100 ecological questions of high policy relevance in the UK." *Journal of Applied Ecology* **43**: 617-627.
- Sutton, R. T., and Hodson, D.L.R. (2005). "Atlantic Ocean Forcing of North American and European Summer Climate." *Science* **309**: 115-117.
- Terray, L., M.E., Demory, M. Déqué, G. de Coetlogon and E. Maisonave. (2004). "Simulation of late-21st-century changes in wintertime atmospheric circulation over Europe due to anthropogenic causes." *Journal of Climate*. **17**: 4630-4635.
- Thomas, C. D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, I.J., Collingham, Y.C., Erasmus, B.F.N., Ferreira de Siqueira, M., Granger, A., Hannah, L., Hughes, L., Huntley, B., Van Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Peterson, A.T., Philips, O.L., and Williams, S.E. (2004). "Extinction risk from climate change." *Nature* **427**: 145-148.
- Thompson, P., and Ollason, J.C. (2001). "Lagged effects of ocean climate change on fulmar population dynamics." *Nature* **413**: 417-420.

- Thuiller, W., Lavorel, S., Araujo, M.B., Sykes, M.T., and Prentice, I.C. (2005). "Climate change threats to plant diversity in Europe." *Proceedings of the National Academy of Sciences* **102**: 8245-8250.
- Thoreau, H. D. (1854). *Walden; or, Life in the Woods*. Boston, Ticknor and Fields.
- Thuiller, W., Albert, C., Araujo, M.B., Berry, P.M., Cabeza, M., Guisan, A., Hickler, T., Midgley, G.F., Paterson, J., Schurr, F.M., Sykes, M.T., Zimmerman, N.E. (2008). "Predicting global change impacts on plant species' distributions: Future challenges." *Perspectives in Plant Ecology, Evolution and Systematics* **9**: 137-152.
- Thuiller, W., Lavorel, S., Araujo, M.B., Sykes, M.T., and Prentice, I.C. (2005). "Climate change threats to plant diversity in Europe." *Proceedings of the National Academy of Sciences* **102**: 8245-8250.
- Tischendorf, L., and Fahrig, L. (2000). "On the usage and measurement of landscape connectivity." *Oikos* **90**: 7-19.
- UCC (2008). PLANFORBIO News. Newsletter, Issue 3, June 2008. **3**: pp4.
- Van der Sluijs, J. P. (1997). *Anchoring Amid Uncertainty: on the Management of of Uncertainties in Risk Assessment of Anthropogenic Climate Change*. Utrecht University, Netherlands.
- Van der Sluijs, J. P., Van Eijdhoven, J., Shackley, S., Wynne, B. (1998). "Anchoring devices in science for policy: The case of consensus around climate sensitivity." *Social Studies of Science* **28**: 291-323.
- Van Vuuren, D. P., O'Neill, B.C. (2006). "The consistency of IPCC's SRES Scenarios to recent literature and recent projections." *Climatic Change* **75**: 1-2.
- Visser, M. E., Holleman, L.J.M. (2001). "Warmer springs disrupt the synchrony of oak and winter moth phenology." *Proceedings of the Royal Society, London B*, **268**: 289-294.
- Walmsley, C. A., Smithers, R.J., Berry, P.M., Harley, M., Stevenson, M.J., Catchpole, R. (Eds.) (2007). MONARCH -Modelling Natural Resource Responses to Climate Change - a synthesis for biodiversity. Oxford, UKCIP: 100pp.
- Walther, B. R., Beissner, S., Burga, C.A. (2005). "Trends in the upward shift of alpine plants." *Journal of Vegetation Science* **16** (5): 541-548.
- Wanless, S., Wright, P.J., and White, S. (2004). "Evidence for decrease in size of lesser sandeels *Ammodytes marinus* in a North Sea aggregation over a 30-year period." *Marine Ecology Progress Series* **294**: 237-246.
- Wanless, S., Harris, M.P., Redman, P. and Speakman, J.R. (2005). "Low energy values of fish as a probable cause of a major seabird breeding failure in the North Sea." *Marine Ecology Progress Series* **294**: 1-8.
- Watts, K., Humphrey, J.W., Griffiths, C., Quine, C., and Ray, D. (2005). Evaluating Biodiversity in Fragmented Landscapes: Principles. Edinburgh, Forest Research Information Note 073: pp8.
- Watts, K., Ray, D., Quine, C., Humphrey, J.W., and Griffiths, M. (2007). Evaluating Biodiversity in Fragmented Landscapes: Applications of Landscape Ecology Tools. Edinburgh, Forest Research Information Note 085: pp8.

- White, M. A., Brunsell, N., Schwartz, M.D. (2003). Vegetation phenology in global change studies. Phenology: An Integrative Environmental Science. M. D. E. Schwartz. Dordrecht, Kluwer Academic Publishers: pp 453-466.
- Williams, P. H., Hannah, L., Andelman, S., Midgley, G.F., Araujo, M.B., Hughes, G.O., Manne, L., Martinez-Meyer, E., Pearson, R.G. (2005). "Planning for climate change: identifying minimum-dispersal corridors for the Cape Proteaceae." *Conservation Biology* **19**: 1063-1074.
- Willows, R., Connell, R. (Eds.) (2003). Climate adaptation: Risk, uncertainty and decision making. Oxford, United Kingdom Climate Impacts Programme Technical Report.
- Woolf, D., and Coll, J. (2006). Impacts of Climate Change on Storms and Waves. Marine Climate Change Impacts Annual Report Card 2006. P. J. Buckley, Dye, S.R. and Baxter, J.M. (Eds.). Lowestoft, MCCIP, Lowestoft, www.mccip.org.uk. Report: pp154.
- Woolf, D., and Coll, J. (2007). Impacts of Climate Change on Storms and Waves. Marine Climate Change Impacts Annual Report Card 2007. P. J. Buckley, Dye, S.R. and Baxter, J.M. (Eds.). Lowestoft, MCCIP, Lowestoft, www.mccip.org.uk.
- Wyse-Jackson, P. (2008). The potential impact of climate change on native plant diversity in Ireland. Short Comm. National Botanic Gardens, Glasnevin. www.botanicgardens.ie/news/20080122.htm, 22 January. Dublin: pp8.
- Zavaleta, E. S., Shaw, M.R., Chiariello, N.R., Thomas, B.D., Cleland, E.E., Field, C.B., Mooney, H.A. (2003). "Grassland responses to three years of elevated temperature, CO₂, precipitation and N deposition." *Ecological Monographs* **73**: 585-604

Appendix 1: Glossary of terms

Adaptation: Adjustment in natural or human systems to a new or changing environment <http://glossary.eea.europa.eu/EEAGlossary>

Climate change context: Initiatives and measures to reduce the vulnerability of natural and human systems against actual or expected climate change effects. Various types of adaptation exist, e.g. anticipatory and reactive, private and public, and autonomous and planned. Examples are raising river or coastal dikes, the substitution of more temperature- shock resistant plants for sensitive ones, etc. IPCC Fourth Assessment Report Synthesis, 2007

Policy context): Policies, practices and projects which can either moderate damage and/or realize opportunities associated with climate change Piper et al., 2006

Evolutionary context: Characteristics of organisms evolved as a consequence of natural selection in its evolutionary past which result in a close match with features of the environment. Begon, Harper & Townsend 1996

Barrier: A physical, chemical or biological feature between two landscape elements that prevents the flow of individuals, genetic material, energy etc. Forman & Godron 1986.

Biodiversity/Biological diversity: The variability among living organisms from all sources including, *inter alia*, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystems. Convention on Biological Diversity:<http://www.cbd.int/convention/articles.shtml?a=cbd-02>.

Buffer zone: The region near the border of a protected area; a transition zone between areas managed for different objectives. <http://www.unepwcmc.org/reception/glossaryA-E.htm>.

Zone / area around the network (ie around core areas and, if necessary, around linkage elements) which protects the network from potentially damaging external influences and which are essentially transitional areas characterized by compatible land uses. Kettunen et al., 2007 (IEEP glossary).

Catchment/catchment area: (1) An area from which surface runoff is carried away by a single drainage system. (2) The area of land bounded by watersheds draining into a river, basin or reservoir. http://glossary.eea.europa.eu/EEAGlossary/C/catchment_area.

Climate change: A change of climate which is attributed directly or indirectly to human activity that alters the composition of the global *atmosphere* and which is in addition to natural climate variability observed over comparable time periods'. *United Nations Framework Convention on Climate Change (UNFCCC)*

Climate change refers to a change in the state of the climate that can be identified (e.g., by using statistical tests) by changes in the mean and/or the variability of its properties, and that persists for an extended period, typically decades or longer. Climate change may be due to natural internal processes or external forcings, or to persistent anthropogenic changes in the composition of the atmosphere or in land use. IPCC Fourth Assessment Report Synthesis, 2007

Colonisation: The entry and spread of a species into an area, habitat or population from which it was previously absent. Begon, Harper & Townsend 1996.

The successful establishment of an invading species in a habitat. Oxford Dictionary of Ecology.

Configuration: The location and juxtaposition of different landscape elements. Forman & Godron 1986.

Connectedness /structural connectivity: Structural connectivity is equal to habitat continuity and is measured by analyzing landscape structure, independent of any attributes of organisms. Kettunen *et al.*, 2007 (IEEP glossary).

The physical distance between elements of the same type (e.g. forest patches). Farina 2000.

(Mathematical context) A space is completely connected if it is not divided into two open wholes (i.e. is not crossed by a boundary whose ends join the perimeter of the space). Hocking & Young 1961.

Connectivity: The functional relationship among habitat patches, owing to the spatial contagion of habitat and the movement responses of organisms to the landscape structure. With *et al.*, 1997.

*The degree to which the landscape facilitates or impedes movement among resource patches. Taylor *et al.*, 1993.*

*Functional connectivity is the response of the organism to the landscape elements other than it's habitats (i.e the non-habitat matrix). Kettunen *et al.*, 2007 (IEEP glossary).*

Core areas: The main places within the landscape where the species/habitats that are the target of the network reside – they may support particular plants or animals, vegetation types, habitats, structural or others features, and can vary in terms of their size, composition, condition, seasonal usage, designation, degree of protection, etc. (and may be graded accordingly) – although it is convenient to delimit a core area within a single site boundary, in reality the situation is likely to be more complex because of variation in composition, condition, etc., within the core area and the supporting role played by external features, in supply food to animals that nest/roost within the core area. Various sources.

Corridor: Landscape elements which serve to maintain vital ecological or environmental connections by providing physical ([continuous] though not necessarily linear) linkages between the core areas. Bennett 2004.

(1) A linear strip of land identified for present or future location of transportation or utility rights of-way within its boundaries.

(2) A thin strip of vegetation used by wildlife and potentially allowing movement of biotic factors between two areas . <http://glossary.eea.europa.eu/EEAGlossary/C/corridor>

A narrow strip of land that differs from the matrix on either side. Forman & Godron 1986.

Dispersal: The spreading of individuals away from each other, e.g. of offspring from their parents or from regions of high population density to regions of lower density. Begon, Harper & Townsend 1996.

Animal dispersal – *A one-way movement of an individual from one home range to a new home range*

Plant dispersal – *A process of plant propagule movement that results in establishment of the species at a new site.* Forman & Godron 1986.

Disturbance: A discrete event, either natural or human induced, that causes a change in the existing condition of an ecological system. Kettunen *et al.*, 2007 (IEEP glossary).

an event that causes a significant change from the normal pattern in an ecological system. Forman & Godron 1986.

Ecological coherence: Sufficient representation of habitats and features of the landscape essential for the migration, dispersal and genetic exchange of wild species are maintained Adapted from: <http://www.bfn.de/fileadmin/MDB/documents/themen/natura2000/conclusions.pdf> (Workshop on Article 10 Habitats Directive 9-12 May 2005 – Conclusions).

Such features are those which, by virtue of their linear and continuous structure (such as rivers with their banks or the traditional systems for marking field boundaries) or their function as stepping stones (such as ponds or small woods), are essential for the migration, dispersal and genetic exchange of wild species. European Habitats Directive article 10.

Sufficient representation of habitats / species to ensure favourable conservation status of habitats and species across their whole natural range. 'Sufficient representation' is a function of patch quality, total patch area, patch configuration and landscape permeability. Kettunen *et al.*, 2007 (IEEP glossary).

Edge effect: A different species composition, relative abundance, species interaction or fitness in the outer part of a patch compared to the interior. Ries *et al.*, 2004.

The existence of more species in a region of overlap between two ecosystems than occur in either of those systems. Oxford Dictionary of Ecology.

Effect size (In meta-analysis): A statistical standardisation of study findings, in a way that means resulting values are interpretable in a consistent fashion across all the variables and measurements involved. Lipsey & Wilson 2001.

Element: The smallest spatial units of ecological community that make up a landscape (habitat patches, matrix patches, barriers, corridors, etc). Forman & Godron 1986.

Emigration: The movement of individuals out of a population or from one area to another. Begon, Harper & Townsend 1996.

Establishment: The arrival of immigrant individuals and subsequent successful life cycle of a viable population. Various sources.

Fragmentation: The breaking-up of continuous tracts of ecosystems creating barriers to migration or dispersal of organisms and reducing the size of homogenous areas.

Fragmentation may be induced by human activities (e.g. road infrastructures, dams) or by natural processes. <http://glossary.eea.europa.eu/EEAGlossary/F/fragmentation>

Habitat: The environment in which an animal or plant lives, generally defined in terms of vegetation and physical features <http://www.unepwcmc.org/reception/glossaryF-L.htm>

An association of species that have a strong spatial association that is quantifiable. Various sources.

The place or type of site where an organism or population naturally occurs. <http://www.cbd.int/convention/articles.shtml?a=cbd-02>

Heterogeneity: Landscapes' quality or state of being heterogeneous, e.g. being composed of parts of different habitats. Kettunen *et al.*, 2007 (IEEP glossary).

Differences in the spatial distribution of species, energy and materials.

Immigration: Entry of organisms into a population [or area] from elsewhere. Begon, Harper & Townsend 1996.

Intensive land use: Land use characterised by a high degree of modification or a high level of inputs. Various sources.

Invasive species: Those which cause rapid changes in ecological communities and thereby species loss. Various sources.

Non-native species which threaten ecosystems, habitats or species.
http://glossary.eea.europa.eu/EEAGlossary/I/invasive_species.

Isolation: The [geographical] prevention of random reproduction among individuals of a population, generally leading to genetically different sub-populations. Forman & Godron 1986.

Landscape: The traits, patterns, and structure of a specific geographic area, including its biological composition, its physical environment, and its anthropogenic or social patterns. Various sources.

An area where interacting ecosystems are grouped and repeated in similar form.
<http://glossary.eea.europa.eu/EEAGlossary/L/landscape>.

Links/Linkages: General term for an arrangement of habitat (not necessarily continuous or linear) that enhances the movement of animals or the continuity of species processes through the landscape. Bennett 1999, (IUCN).

Matrix: The interstitial habitat / environment between habitat patches in a habitat mosaic, typically comprising the most extensive habitat /environment type in a landscape. Kettunen *et al.*, 2007 (IEEP glossary).

The most extensive and most connected landscape element type present, which plays the dominant role in landscape functioning. Also, a landscape element surrounding a patch.
Forman & Godron 1986.

Meta-analysis: Statistical aggregation and comparison of a number of empirical studies. Lipsey & Wilson, 2001.

Migration: A cyclic movement of animals between separated areas that are used during different seasons. Forman & Godron 1986.

The movement of individuals, and commonly whole populations from one region to another. Begon, Harper & Townsend 1996.

Mitigation: Technological change and substitution that reduce resource inputs and emissions per unit of output. Although several social, economic and technological policies would produce an emission reduction, with respect to Climate Change, mitigation means implementing policies to reduce greenhouse gas emissions and enhance sinks. IPCC Fourth Assessment Report Synthesis, 2007.

Mosaic (habitat mosaic): Spatial configuration of habitats within a landscape, generally formed by patches arranged within a matrix. Kettunen *et al.*, 2007 (IEEP glossary).

Movement: Any movement by an individual, directed or undirected, permanent or temporary (c.f. dispersal). Various sources.

Network: A coherent system of natural and/or semi-natural landscape elements that is configured and managed with the objective of maintaining or restoring ecological functions as a means to conserve biodiversity while also providing appropriate opportunities for the sustainable use of natural resources. Bennett 2004.

Patch: A particulate, invariant and homogeneous entity within an ecosystem. Kettunen *et al.*, 2007 (IEEP glossary).

A nonlinear surface area differing in appearance from its surroundings. Forman & Godron 1986.

Permeability: The quality of a heterogeneous land area to provide for passage of organisms. Kettunen *et al.*, 2007 (IEEP glossary).

Protected area: An area of land and/or sea especially dedicated to the protection and maintenance of biological diversity, and of natural and associated cultural resources, and managed through legal or other effective means <http://www.unepwcmc.org/reception/glossaryM-R.htm>

Range: The spatial limits within which the habitat or species occurs. A natural range is not static but dynamic: it can decrease and expand. IPCC Fourth Assessment Report Synthesis, 2007/Habitats directive.

Resilience: A tendency to maintain integrity when subject to disturbance. UNDP, 2007.

The ability of a social or ecological system to absorb disturbances while retaining the same basic structure and ways of functioning, the capacity for self-organisation, and the capacity to adapt to stress and change. IPCC Fourth Assessment Report Synthesis, 2007.

The ability of, or speed at which, a community or population returns to its original state after a disturbance, or its ability to resist the impacts of disturbance. EBS (Mitchell *et al.*, 2007).

The magnitude of disturbance that can be absorbed before the system changes its structure by changing the variables and processes that control behaviour. Or: the speed with which a disturbed system returns to equilibrium or the same general state after being changed. Piper *et al.*, 2006.

Resistance (Connectivity context): The inverse of permeability. Various sources.

Restoration: The process of assisting the recovery of an ecosystem that has been degraded, damaged, or destroyed.

http://www.ser.org/content/ecological_restoration_primer.asp#3

Where substantial effort is needed to [restore] a site with relict features (or historically former habitat).

<http://www.ukbap.org.uk/library/brig/TargetsReview06/Final/BAPTtargetDefinitionsGuidance.pdf>

The return of an ecosystem or habitat to its original community structure, natural complement of species, and natural functions. <http://www.unepwcmc.org/reception/glossaryM-R.htm>

Measures taken to return a site to pre-violation conditions.

<http://glossary.eea.europa.eu/EEAGlossary/R/restoration>

Semi-natural (Vegetation/habitat context): Communities consisting predominantly of native species that have not been planted but have arisen from natural regeneration, i.e. from seed or clonal regrowth. Adapted from Spencer & Kirby, 1992

Both natural and semi-natural vegetation may be found in upland and lowland situations, including wetland, woodland, grassland, moor and heathland habitats, and are described as being made up of self-seeded or self propagated vegetation characteristic of the area in which the land is situated. <http://www.crosscompliance.org.uk/faqsGAEC9.htm>

Land that has been subject to past human intervention that has reverted to a more natural state either through recolonisation or expansion of native species. Various sources.

Stepping stone: A patch that allows species to move incrementally across an otherwise hostile landscape (most often associated with migratory [or dispersing] species). Various sources.

A spot that is colonised by a species. Forman & Godron 1986