# **National Parks and Wildlife Service**

## **Conservation Objectives Series**

## Ballyogan Lough SAC 000019





### National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht,

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie E-mail: nature.conservation@ahg.gov.ie

### Citation:

NPWS (2018) Conservation Objectives: Ballyogan Lough SAC 000019. Version 1. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht.

Series Editor: Rebecca Jeffrey ISSN 2009-4086

### Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

### **Notes/Guidelines:**

- 1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
- 2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
- 3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
- 4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
- 5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

### Qualifying Interests

\* indicates a priority habitat under the Habitats Directive

000019 Ballyogan Lough SAC

7210 Calcareous fens with  $\hat{O}|\partial \hat{a} \hat{a} \{\hat{A} \otimes \hat{a} \hat{a} \}$  and species of the Caricion davallianaeE

### Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

#### **NPWS Documents**

**Year:** 1972

Title: A preliminary report on Areas of Scientific Interest in County Clare

Author: Goodwillie, R.N.

Series: Unpublished report

Year: 2012

Title: Ireland Red List No. 8: Bryophytes

Author: Lockhart, N.; Hodgetts, N.; Holyoak, D.

Series: Ireland Red List series, NPWS

Year: 2013

Title: The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments

Author: NPWS

Series: Conservation assessments

Year: 2013

Title: Conservation status assessments for three fen habitat types - 7230, 7210 and 7140

Author: Kimberley, S.

Series: Unpublished report to NPWS

Year: 2014

Title: Guidelines for a national survey and conservation assessment of upland vegetation and

habitats in Ireland, Version 2.0

Author: Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.

Series: Irish Wildlife Manual No. 79

**Year:** 2016

Title: Ireland Red List No. 10: Vascular Plants

Author: Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.;

Wright, M.

Series: Ireland Red Lists series, NPWS

### **Other References**

Year: 2004

Title: Common Standards Monitoring guidance for lowland wetland habitats

Author: JNCC

Series: Joint Nature Conservation Committee, Peterborough

### Conservation Objectives for: Ballyogan Lough SAC [000019]

## 7210 Calcareous fens with Cladium mariscus and species of the Caricion dayallianae

To maintain the favourable conservation condition of Calcareous fens with *Cladium mariscus* and species of the Caricion davallianae\* in Ballyogan Lough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Calcareous fens with <i>Cladium mariscus</i> and species of the Caricion davallianae* has not been mapped in detail for Ballyogan Lough SAC and thus the total area of the qualifying priority habitat in the SAC is unknown. The SAC supports typical examples of the habitat in a calcareous lake system, which includes Ballyogan Lough, Moyree Lough and other small lakes. The habitat occurs in a mosaic with other swamp and fen/marsh wetland habitats and cutover bog vegetation predominantly in the north-east of the SAC. Greater fen-sedge ( <i>Cladium mariscus</i> ) dominated areas mostly occur in the wetter areas close to the lakes (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for Habitat area above
Ecosystem function: peat formation	Percentage cover of peat-forming vegetation and water table levels	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time
Ecosystem function: hydrology - groundwater levels	Water levels (centimetres); duration of levels; hydraulic gradients	Maintain, or where necessary restore, appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Fen habitats require high groundwater levels (i.e. water levels at or above the ground surface) for a large proportion of the calendar year (i.e. duration of mean groundwater level). Fen groundwater levels are controlled by regional groundwater levels in the contributing catchment area (which sustain the hydraulic gradients of the fen groundwater table). Regional abstraction of groundwater may affect fen groundwater levels
Ecosystem function: hydrology - surface water flow	Drain density and form	Maintain, or where necessary restore, as close as possible to natural or semi-natural drainage conditions	Drainage, either within or surrounding the fen habitat, can result in the drawdown of the fen groundwater table. The depth, geometry and density of drainage (hydromorphology) will indicate the scale and impact on fen hydrology. Drainage can result in loss of characteristic species and transition to drier habitats
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly pH and nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources.  However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient under natural conditions. Water supply should also be relatively calcium-rich
Vegetation composition: typical species	Percentage cover at a representative number of 2m x 2m monitoring stops	Maintain adequate cover of typical species, including brown mosses and vascular plants	For lists of typical plant species, see the Article 17 conservation status assessment for <i>Cladium</i> fens (NPWS, 2013) and the Article 17 fen habitats supporting document (Kimberley, 2013). In this SAC, stands of greater fen-sedge ( <i>Cladium mariscus</i> ) occur in association with common reed ( <i>Phragmites australis</i> ), and species including black bog-rush ( <i>Schoenus nigricans</i> ), blunt-flowered rush ( <i>Juncus subnodulosus</i> ) and small sedges such as dioecious sedge ( <i>Carex dioica</i> ) and lesser tussock-sedge ( <i>C. diandra</i> ) (Goodwillie, 1972; NPWS internal files)
Vegetation composition: native negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of native negative indicator species at insignificant levels	Negative indicators include species not characteristic of the habitat and species indicative of undesirable impacts such as overgrazing, undergrazing, nutrient enrichment, agricultural improvement or impacts on hydrology. See JNCC (2004) and Kimberley (2013)

Vegetation composition: non- native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014). Scrub and trees will tend to invade if fen conditions become drier
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground not more than 10%. Where tufa is present, disturbed bare ground not more than 1%	Attribute and target based on Perrin et al. (2014). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes. Disturbance can include hoof marks, wallows, human footprints, vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for peatlands
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat; maintain features of local distinctiveness, subject to natural processes	This includes species on the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

