



## Report of the profile of COVID-19 cases in healthcare workers in Ireland

Report prepared by HPSC on 04/08/2020

## Note: Data are provisional

## **Summary**

Healthcare worker (HCW) status is determined both by self-classification and workplace. The definition includes anyone who self- identifies as a HCW irrespective of where they work. In addition, all staff that work in any healthcare facility (includes cleaners, household staff etc.) are classified as HCWs. The category includes healthcare workers employed both by public and private providers.

This report includes data as of 4<sup>h</sup> August 2020 at 9:15 am for events created on CIDR up to midnight Saturday 1<sup>st</sup> August 2020.

Characteristic of HCW COVID-19 cases	Number	%
Total number of COVID-19 cases	26628	-
Total number of HCW cases	8500	31.9
Number of new cases reported between 26 July – 1August	19	-
Median age (IQR)	41 (31-50) years	-
Total number hospitalised	327	3.8
The median age (IQR) of hospital inpatient	46 (34-53) years	
Total number admitted to ICU	53	16.2*
The median age (IQR) of HCW in ICU	51 (43-60) years	-
Total number of deaths	8**	0.09
The median age (range) for deaths	54 (30-68) years	-

<sup>\*</sup>This relates to hospitalised cases and it is 0.6% of all HCW cases.

There was a total of 26628 COVID-19 cases reported as of 01/08/2020 12.00 midnight, 8500 (31.9%) were healthcare workers (HCWs).

The median age of COVID-19 case HCWs is 41 years (range 17-81 years). The proportion of females is disproportionately high (74.0%) among HCW COVID-19 cases compared to non-HCW COVID-19 cases (49.0%), most likely due to some HCW specialities e.g. nursing, being female dominated. The majority (58.3%) of notified HCWs were from the HSE East. CIDR (Computerised Infectious Diseases Reporting) is a dynamic information system and data are continuously validated and updated.

<sup>\*\*</sup>Seven confirmed and 1 probable COVID-19 case.

This report includes all HCW COVID-19 cases (n=8500) reported to HPSC, including confirmed, probable and possible cases. There are 63 (0.7%) probable or possible HCW COVID-19 cases included in the report.

Table 1. Number and proportion of HCW COVID-19 cases by epidemiological week

Epi Week (Calendar Date)	Number of HCWs	Total number of cases*	Proportion of HCWs (%)
10 (1 – 7, March)	3	16	18.8
11 (8 – 14, March)	23	128	18.0
12 (15 – 21, March)	205	683	30.0
13 (22 – 28, March)	441	1631	27.0
14 (29 March - 4 April)	945	2433	38.8
15 (5 – 11, April)	1631	4580	35.6
16 (12 – 18, April)	1624	5664	28.7
17 (19 - 25, April)	1252	3892	32.2
18 (26 April – 2 May)	825	2583	31.9
19 (3 –9 May)	501	1500	33.4
20 (10-16 May)	487	1187	41.0
21 (17-23 May)	149	585	25.5
22 (24-30 May)	106	425	24.9
23 (31 May - 6 June)	73	243	30.0
24 (7 – 13 June)	23	114	20.2
25 (14-20 June)	29	102	28.4
26 (21-27 June)	18	63	28.6
27 (28 June- 4 July)	25	94	26.6
28 (5-11 July)	43	139	30.9
29 (12- 18 July)	45	154	29.2
30 (19-25 July)	33	124	26.6
31 (26 July – 1 August)	19	288	6.6
Total	8500	26628	31.9

<sup>\*</sup>including HCWs; confirmed, probable and possible cases.

Figure 1. Proportion of HCW and non-HCW COVID-19 cases by week in which case was notified (n= 26628)






Figure 2. Distribution of HCW COVID-19 cases by week in which case was notified \* and age group (n=8491)



<sup>\*</sup> Age was not available for 9 cases.

Figure 3. Distribution of HCW COVID-19 cases by age and sex (n=8476) \*



<sup>\*</sup> Age and sex were not available for 24 cases.

Table 2. Number and proportion of HCW COVID-19 cases by HSE area

HSE area	Number of HCWs cases	Proportion of all HCW cases	Number of new HCW cases since last week
HSE E	4956	58.3%	13
HSE M	453	5.3%	0
HSE MW	369	4.3%	<5
HSE NE	1172	13.8%	<5
HSE NW	309	3.6%	<5
HSE SE	349	4.1%	0
HSE S	460	5.4%	<5
HSE W	432	5.1%	<5
Total	8500	100.0%	19

Table 3. Number and proportion of HCW COVID-19 cases by CHO area

CHO area	Number of HCWs cases	Proportion of all HCW cases	Number of new HCW cases since last week
CHO1	828	9.7%	<5
CHO2	432	5.1%	<5
CHO3	369	4.3%	<5
CHO4	460	5.4%	<5
CHO5	349	4.1%	0
CHO6	916	10.8%	<5
CHO7	1822	21.5%	5
CHO8	1106	13.0%	<5
CHO9	2218	26.1%	7
Total	8500	100.0%	19

Table 4. Number and proportion of HCW COVID-19 cases by role

HCW Role	Number of HCWs cases	Proportion of all HCW cases	Number of new HCW cases since last week
Nurse	2746	32.3%	<5
Healthcare assistant	2261	26.7%	<5
Doctor	524	6.1%	0
Porter	97	1.1%	<5
Other HCW	1995	23.5%	7
Unknown	877	10.4%	5
Total	8500	100.0%	19

Table 5. Number and proportion of HCW COVID-19 cases linked to an outbreak by outbreak location

Outbreak location	Number of HCWs cases	Proportion of all HCW cases	Number of new HCW cases since last week
Nursing home	2096	24.8%	0
Hospital	727	8.5%	0
Private house	688	8.1%	<5
Residential institution	444	5.2%	0
Comm. Hosp/Long-stay unit	233	2.7%	<5
Workplace	33	0.4%	0
Travel related	30	0.3%	<5
Extended family	17	0.2%	0
Community outbreak	14	0.2%	0
Public house	2	0.0%	0
Restaurant / Cafe	1	0.0%	0
Hotel	1	0.0%	0
Unknown	1	0.0%	0
Not Specified	1	0.0%	0
Other	22	0.3%	0
Not linked to an outbreak	4190	49.3%	16
Total	8500	100.0%	19

Table 6. Number and proportion of HCW COVID-19 cases with underlying medical conditions

Underlying clinical conditions	Number	Percentage
Yes	2957	34.8%
No	4735	55.7%
Unknown	808	9.5%
Total	8500	100.0%

Table 7. Number of confirmed COVID-19 cases by WHO transmission classification\*

Transmission classification*	Number	Percentage
Community transmission - including possible community transmission**	1180	13.9%
Local transmission	7173	84.4%
Travel related	147	1.7%
Total	8500	100.0%

<sup>\*</sup>WHO definition of transmission classification is specified below:

'Most likely source of transmission' is a composite variable created by combining several data fields on CIDR. The rate of HCWs with 'under investigation' transmission source category (n=865 10.1%) is expected to decrease due to continuous improvement of the transmission source variable algorithm and the completeness of surveillance data.

Table 8. Most likely source of transmission of COVID-19 in HCWs

Likely source of transmission	Number	Percentage
Healthcare setting acquired: staff*	6198	73.0%
Close contact with a known confirmed case	962	11.2%
Travel related	147	1.7%
Community transmission	315	3.7%
Healthcare setting acquired: patient	13	0.2%
Under investigation	865	10.1%
Total	8500	100.0%

\*Includes HCWs with most likely source of transmission notified as 'Healthcare setting acquired: staff' AND HCWs with most likely source of transmission field not completed on CIDR who had close contact with a COVID-19 case in healthcare or workplace setting AND HCWs with most likely source of transmission field not completed on CIDR who are linked to an outbreak in a Comm. Hosp/Long-stay unit / Hospital / Nursing home.

<sup>-</sup> Community transmission is evidenced by the inability to relate confirmed cases through chains of transmission for a large number of cases, or by increasing positive tests through routine screening of sentinel samples.

<sup>-</sup> Local transmission indicates locations where the source of infection is within the reporting location.

<sup>-</sup> Imported cases only indicates locations where all cases have been acquired outside the location of reporting.

<sup>\*\*</sup> Community transmission relates to those in table 11 below who are in the two categories community transmission and under investigation.

## Acknowledgements

Sincere thanks are extended to all those who are participating in the collection of data and reporting of data used in these reports. This includes the HSE COVID-19 Contact Management Programme (CMP), staff in ICU units, notifying clinicians, laboratory staff, public health doctors, nurses, surveillance scientists, microbiologists and administrative staff.