

## THE SYSTEMIC REDSHIFT OF THE QUASAR 1331 + 170

R. F. CARSWELL

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, England

C. M. MOUNTAIN, D. J. ROBERTSON, S. M. BEARD, A. R. GLENDINNING, D. C. LAIRD, L. C. LAWRENCE,  
 D. MONTGOMERY, G. PENTLAND, D. A. PICKUP, AND I. A. SMITH

Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, Scotland

J. A. BAILEY, A. BRIDGER, M. M. CASALI, T. R. GEBALLE, P. PUXLEY, M. G. SMITH, AND G. S. WRIGHT  
 Joint Astronomy Centre, 665 Komohana Street, Hilo, HI 96720

S. K. RAMSAY

Department of Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, Scotland

A. C. BAKER

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, England

B. R. ESPEY

Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260

AND

M. J. WARD

Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, England

Received 1991 June 18; accepted 1991 August 14

### ABSTRACT

Infrared spectra covering the 1.6 and 2  $\mu\text{m}$  regions of the quasar 1331 + 170, redshift  $z = 2.09$ , were obtained with the CGS4 spectrometer at the UK Infrared Telescope. These spectra show redshifted  $\text{H}\alpha$ ,  $\text{H}\beta$ , and  $[\text{O III}]$  lines. The redshifts of the forbidden lines are similar to the low-ionization lines measured at optical wavelengths but are significantly higher than the high-ionization line redshift. Thus the low-ionization lines are likely to provide a reliable estimate for the systematic redshift in this object, and the high-ionization material is therefore flowing out from, or into, the quasar and is subject to some obscuration.

*Subject headings:* line profiles — quasars

### 1. INTRODUCTION

Work by Gaskell (1982) and Wilkes (1984) showed that there are systematic profile and velocity differences between the ultraviolet emission lines seen shifted to optical wavelengths in the spectra of high-redshift quasars. Subsequent detailed studies (Espey et al. 1989; Corbin 1990) have confirmed this finding. For example, Espey et al. measured line profiles for  $\text{Ly}\alpha$ ,  $\text{C IV } \lambda 1549$ ,  $\text{C III } \lambda 1909$ ,  $\text{Mg II } \lambda 2798$ , and  $\text{H}\alpha$  in a number of quasars. They showed that for their sample the high-ionization lines and  $\text{Ly}\alpha$  have lower redshifts by around  $1000 \text{ km s}^{-1}$  compared to the low-ionization lines and  $\text{H}\alpha$ . A number of possible explanations for these velocity differences have been suggested. They generally involve bulk mass flows and internal obscuration in the quasar, but since the true redshifts of the quasars were not known, it has been difficult to choose between them. One obvious way forward is to measure the forbidden line redshifts. Since these are expected to arise in the extended emission regions, they should therefore provide a reasonably accurate value of the systemic redshift for the quasar.

There have been previous attempts to measure the forbidden lines in high-redshift quasars, but these have not been of sufficient accuracy to permit useful estimates of the redshift. Kühr et al. (1984) report the detection of the  $[\text{O III}] \lambda\lambda 4959, 5007$  lines in the  $z = 3.4$  quasar 0014 + 81, but in this case all lines have the same redshift to within the errors. Espey et al. (1989) attempted to measure the same lines in the quasar 1100 – 264, at a redshift of  $z = 2.14$ , in which there is a significant velocity

shift between ionization levels, but they failed to attain sufficient signal-to-noise ratio in their spectrum to detect the lines.

We report here the measurement of Balmer line and  $[\text{O III}]$  emission shifted to infrared wavelengths in the spectrum of the quasar 1331 + 170 which has a redshift of 2.09. This quasar was first discovered as part of a radio source identification program (Baldwin, Wampler, & Gaskell 1973) and was chosen for this study because it is known to have a large velocity shift between low- and high-ionization lines (Espey et al. 1989). It thus provides a straightforward example of the profile differences between the various broad emission lines.

### 2. OBSERVATIONS

A spectrum of 1331 + 170 covering the  $H$  window was obtained using the Cooled Grating Spectrometer CGS4 (Mountain et al. 1990) during its first scheduled commissioning run on UKIRT on 1991 February 11, and subsequent spectra in both  $H$  and  $K$  were obtained in 1991 April. The detector for all observations was a SBRC  $62 \times 58$  InSb array. Since the pixel size of  $76 \mu\text{m}$  corresponds to  $3''$ , to adequately sample the instrument profile, the detector was moved in the spectral direction between exposures. The first observation consisted of 24 object/sky pairs at three detector positions each with an exposure time of 20 s. The total integration time on the source was 24 minutes, with an equal time devoted to sky observations. For subsequent observations the object was switched between two defined positions along the slit, and the remainder of the slit was used to measure the sky. During all the  $H$ -band

TABLE 1  
OBSERVATIONS

Date (1991)	Exposure (minutes)	Grating (lines mm <sup>-1</sup> )	Order	Wavelength Range (μm)	Substeps per Pixel	Resolution (λ/δλ)	Comments
Feb 11 .....	24	75	1	1.41–1.82	3	200	Plus 24 minutes of sky observation
Apr 7 .....	115	75	2	1.43–1.64	2	450	Efficiency low
Apr 21 .....	27	75	1	1.95–2.34	3	~300	

exposures, the array was read out using a multiple nondestructive scheme (Chapman et al. 1990), resulting in an effective read noise of 30 electrons per exposure. This, combined with a dark current plus background seen by the array of less than 10 electrons per second per pixel, meant that all the observations were background noise-limited. Details of the observations are given in Table 1.

To extract the object spectrum, the sum of the object minus sky frames was formed, and then an optimal extraction performed using the method described by Horne (1986). The wavelength scales were determined from argon lamp exposures combined with the positions of sky emission lines present in the object exposures.

To determine the [O III] redshift  $z = 2.095$  and line fluxes, we have used the fact that the [O III]  $\lambda 5007$ : [O III]  $\lambda 4959$  flux

ratio is 3:1 and fitted single Gaussian profiles to the lines in the H $\beta$ /[O III] blend. The measured forbidden line widths are close to those of the comparison arc lines, so we are unable to comment on the intrinsic FWHM of the [O III] lines, apart from quoting an upper limit of around 1500 km s<sup>-1</sup>. The other parameters for the emission lines are contained in Table 2.

The 2 μm spectrum shows strong broad H $\alpha$  emission and no other discernible features (Fig. 1). An arc calibration was not taken for this spectrum, so the wavelengths were determined from the sky features. For this reason we are unable to determine the spectral resolution directly, but use the known spectrograph characteristics to give an estimate in Table 1. The redshift given in Table 2 is the result of a Gaussian fit to the line. Against the assumed continuum, the formal uncertainty in the velocity is ~100 km s<sup>-1</sup>, but the result also depends on the




FIG. 1.—The CGS4 spectra of 1331 + 170. The vertical bars indicate the  $\pm 1 \sigma$  error range at each data value. Redshifted H $\alpha$  is at 2.035  $\mu\text{m}$ , H $\beta$  at 1.505  $\mu\text{m}$ , and [O III]  $\lambda 5007$  at 1.545  $\mu\text{m}$ . The fainter continuous curves show the fitted profiles.

continuum placement and slope. Trials with different subsets of the data suggest that the typical uncertainty in the velocity measure is about 200–300 km s<sup>-1</sup>.

### 3. DISCUSSION

The *H*-window spectrum shows the redshifted H $\beta$  and [O III]  $\lambda\lambda$ 4959, 5007 emission lines at redshift  $z = 2.095$  (Fig. 1), and an otherwise featureless continuum. The spectral resolution is sufficiently high that the lines should be clearly separated, but at the signal-to-noise ratio achieved the [O III]  $\lambda$ 4959 line is poorly defined. We believe that the lines we have identified as [O III] are predominantly [O III] and not Fe II emission from multiplet 42 at wavelengths 4924 and 5018 Å (in contrast with 3C 273; Shuder 1984). This is because the feature is clearly narrower than H $\beta$  and there is no sign of the other component of Fe II multiplet 42, Fe II  $\lambda$ 5169. Also, Fe II emission is generally quite weak in this object—Fe II emission from multiplets 48 and 49 around 5300 Å is not seen in our spectra, and the spectra used by Espey et al. (1989) do not show particularly strong Fe II emission in the region below the Mg II  $\lambda$ 2798 emission line.

The [O III]  $\lambda\lambda$ 4959 + 5077/H $\beta$  intensity ratio is  $\sim 0.75$ . This is similar to the ratio  $\sim 0.85$  found from CVF spectra of the quasar 0014+81 at a redshift of  $z = 3.4$  (Kühr et al. 1984) and is within the range found for lower redshift lower luminosity quasars (see, e.g., Baldwin 1975; Steiner 1981). If these values are typical of bright high-redshift quasars, then the low-redshift example, 3C 273, with [O III]/H $\beta \sim 0.12$  (Shuder 1984), is therefore atypical of the class.

The CGS4 data presented here supplement the line profile measurements obtained by Espey et al. (1989) for Ly $\alpha$ –Mg II  $\lambda$ 2798 and H $\alpha$ . Espey et al. found that the high-ionization lines (and Ly $\alpha$ ) are blueshifted with respect to the low-ionization lines (plus H $\alpha$ ) in 1331+170 and other high-redshift quasars. In the case of 1331+170, the velocity shift is about 1000 km s<sup>-1</sup>, and the dilemma from the earlier study was to identify which lines should be used to determine the true redshift. Since the forbidden lines arise in extended regions beyond the broad-line region where the velocities are considerably lower, they should provide a means of determining the redshift without such ambiguities. Thus the redshift determined from [O III]  $\lambda$ 5007,  $z = 2.095$ , is the true (systemic) redshift of 1331+170. This agrees well with the redshift of Mg II  $\lambda$ 2798 and (less certainly) O I  $\lambda$ 1305 from Espey et al., so the low-ionization lines have a mean redshift close to that of the quasar (see Table 2). The emission-line profiles are shown in Figure 2.

It is clear that these velocity differences require the presence of some form of bulk gas motions, which are different for the various lines concerned. The direction of the flow remains ambiguous. If the obscuration is external to the emission clouds (e.g., as for an accretion disk or distributed continuous absorption near the central continuum source), then the material responsible for the high-ionization emission lines must be flowing out from the quasar. Infall models with electron scattering (Kallman & Krolik 1986), or obscuring material associated with the emission clouds, could equally well explain the observational data.

Our conclusion that the high-ionization material is in bulk flow and suffering obscuration depends on the assumption that the system redshift is given by [O III]. A study of Seyfert galaxies by Whittle (1991) shows that the [O III] peak velocity is generally a good indicator of the systemic velocity, so our assumption appears reasonable. It is possible, however, that in




FIG. 2.—Profiles of the emission lines in 1331+170, reduced to a velocity scale relative to a redshift of  $z = 2.095$ . The Mg II  $\lambda$ 2798 and shorter wavelength lines are from Espey et al. (1989). The Ly $\alpha$  and N V  $\lambda$ 1240 lines are blended in the original spectrum, so for the purposes of illustration they have been separated using the C IV  $\lambda$ 1549 line as a template. The Ly $\alpha$  profile shown is the result of subtracting  $0.9 \times$  the C IV  $\lambda$ 1549 line profile shifted to N V  $\lambda$ 1240, and the N V  $\lambda$ 1240 profile is the result of subtracting  $1.8 \times$  the C IV  $\lambda$ 1549 shifted to Ly $\alpha$ .



TABLE 2  
EMISSION-LINE PARAMETERS

Line	$z$	EW (Å) (rest)	FWHM (km s <sup>-1</sup> )	Velocity cf. [O III]	Observation Date(s)
H $\alpha$ .....	2.101	250	4600	600	1991 Apr 21
[O III] $\lambda$ 5007 .....	2.095	20	1100	...	1991 Feb 11 and Apr 7
[O III] $\lambda$ 4959 .....	2.095	6	1100	...	1991 Feb 11 and Apr 7
H $\beta$ .....	2.097	36	5700	250	1991 Feb 11 and Apr 7
H $\alpha$ .....	2.103	140	3000	800	1987 Mar 16 and 1988 May 1
Mg II $\lambda$ 2798 .....	2.094	16	3200	-100	1988 May 26
C III] $\lambda$ 1909 .....	2.085	22	6600	-1000	1988 May 26
C IV $\lambda$ 1549 .....	2.077	19	4500	-1800	1988 May 26
O I $\lambda$ 1305 .....	2.095	4	3500	0	1988 May 26
N V $\lambda$ 1240 .....	2.080	...	4500	-1500	1988 May 26
Ly $\alpha$ .....	2.083	...	4800	-1200	1988 May 26

an individual case the [O III] emission may arise from material which is itself in bulk motion as, for example, in the Seyfert galaxy Markarian 78 (de Robertis 1987).

The H $\alpha$  line flux is  $\sim 6.5 \times 10^{-17} \text{ W m}^{-2}$ , and that of H $\beta$  is  $1.2 \times 10^{-17} \text{ W m}^{-2}$ , so the H $\alpha$ /H $\beta$  ratio is  $\sim 5.5$ . This value is uncertain, since the spectra were obtained through a 3" slit, so the photometric accuracy may not be very high. Indeed, the sections of continuum observed are flat and suggest that a constant continuum level  $\sim 1 \text{ mJy}$  over the wavelength range might be appropriate. Using a normalization to make the continua levels match, the Balmer line ratio is  $\sim 4.2$ . Also, we have not allowed for the possible presence of [N II]  $\lambda\lambda 6548, 6584$  emission as a contaminant of the H $\alpha$  line. It is difficult to do this unambiguously, but a best fit using the [O III] parameters as templates suggests that about 33% of the feature could be due to forbidden [N II] and [S II]. Thus the H $\alpha$ /H $\beta$  flux ratio is likely to be lower than 5.5 and is probably between 3.7 and 5.5. This is within the range found for many lower redshift quasars and active galactic nuclei (e.g. Baldwin 1975; Rudy 1984; Baldwin et al. 1989).

Comparison of the H $\alpha$  line in the 1991 CGS4 spectrum with that obtained using FIGS at the AAT in 1987 and 1988 reveals some interesting differences. The line equivalent width has increased since the earlier spectra by a factor of about 1.8. The absolute fluxes for these measurements are uncertain since the conditions were not necessarily photometric, but it appears that the continuum flux has decreased by a factor of 2 since the earlier observations. This could naturally account for the change in the H $\alpha$  line equivalent width.

However, there is evidence that the H $\alpha$  line has also varied, since there is a large discrepancy between the H $\alpha$  line profile widths at the two epochs. The 1991 data give H $\alpha$  FWHM 4500 km s<sup>-1</sup>, while the 1987–1988 material yields 3000 km s<sup>-1</sup>. It is hard to see how this difference could arise because of differences in the instrument resolution ( $\sim 800 \text{ km s}^{-1}$  for both the CGS4 and the FIGS data), or as a result of the wings of the line

being lost in the continuum, so the most plausible explanation is an intrinsic variation in the H $\alpha$  profile. Comparison of the optical data from 1987–1988 with those published by Baldwin et al. (1973) shows that any changes in the high-ionization line widths or equivalent widths over 15 yr have been small.

#### 4. CONCLUSIONS

The H $\alpha$ /H $\beta$  flux ratio in 1331+170 has a value  $\sim 3.5$ –5.5, which is similar to that observed in low-redshift quasars. The H $\beta$  line is broader than H $\alpha$ ; this is a common property of active galaxies and low-redshift quasars (Osterbrock & Shuder 1982), and these lines also have a slightly different redshift. The profile width of the H $\alpha$  line seems to have increased significantly, and the continuum flux decreased, over a time scale  $\sim 1$  yr in the quasar's frame, indicating a characteristic size for the Balmer line-emitting region  $\lesssim 1 \text{ lt-yr}$ .

We have detected [O III]  $\lambda$ 5007 at a redshift of 2.095. Comparison with other lines in this object shows that the low-ionization lines have similar redshifts, while the high-ionization material appears to have a systematic velocity toward us with respect to the low-ionization material. If this is true for all quasars which show similar velocity shifts and assuming the forbidden lines are at the systemic redshift, then the high-resolution material must be in ordered motion and suffer some obscuration to explain the differential velocities.

Observations of the forbidden lines in a sample of objects are required to establish reliably which of the broad lines are good indicators of the systemic redshift. Using the new generation of infrared spectrographs, such as CGS4, these observations are now feasible for quasars with redshifts around 2.

We are grateful to the UKIRT telescope operators, Joel Aycock and Thor Wold, for their usual expert assistance and their patience and good humor in coping with a large instrument commissioning team. We also thank Paul Hewett for discussion on the interpretation of the data.

#### REFERENCES

- Baldwin, J. A. 1975, ApJ, 201, 26  
 Baldwin, J. A., Burbidge, E. M., Hazard, C., Murdoch, H. S., Robinson, L. B., and Wampler, E. J. 1973, ApJ, 185, 739  
 Baldwin, J. A., Wampler, E. J., & Gaskell, C. M. 1989, ApJ, 338, 630  
 Chapman, A. R., Beard, S. M., Mountain, C. M., Pettie, D. G., & Pickup, D. A. 1990, Instrumentation in Astronomy VII, ed. D. L. Crawford (Proc. SPIE, 1235), 34  
 Corbin, M. R. 1990, ApJ, 357, 346  
 de Robertis, M. M. 1987, ApJ, 316, 597  
 Espey, B. R., Carswell, R. F., Bailey, J. A., Smith, M. G., & Ward, M. J. 1989, ApJ, 342, 666  
 Gaskell, C. M. 1982, ApJ, 263, 79  
 Horne, K. 1986, PASP, 98, 609  
 Kallman, T. R., & Krolik, J. H. 1986, ApJ, 308, 805  
 Kühn, H., McAlary, C. W., Rudy, R. J., Strittmatter, P. A., & Reike, G. H. 1984, ApJ, 284, L5  
 Mountain, C. M., Robertson, D. J., Lee, T. J., & Wade R. 1990, Instrumentation in Astronomy VII, ed. D. L. Crawford (Proc. SPIE, 1235), 25  
 Osterbrock, D. E., & Shuder, J. M. 1982, ApJS, 49, 149  
 Rudy, R. J. 1984, ApJ, 284, 33  
 Shuder, J. M. 1984, ApJ, 280, 491  
 Steiner, J. E. 1981, ApJ, 250, 469  
 Whittle, D. M. 1991, ApJ, submitted  
 Wilkes, B. J. 1984, MNRAS, 207, 73