



Isolation quick guide -Adults and children from their 13th birthday

V3.4. 16.09.2021

The most common symptoms of COVID-19 are

- Fever
- Dry cough
- Fatigue

Other symptoms that are less common and may affect some patients include:

- Loss of taste or smell,
- Nasal congestion,
- Conjunctivitis (also known as red eyes)
- Sore throat,
- Headache,
- Muscle or joint pain,
- Different types of skin rash,
- Nausea or vomiting,
- Diarrhea,
- Chills or dizziness.

Symptoms of severe COVID-19 disease may include shortness of breath, loss of appetite, confusion, persistent pain or pressure in the chest, high temperature (above 38 °C).

Symptoms may present differently in different age groups and with different variants of the virus. Some symptoms may also be confused with common cold, hay-fever or flu. If you have any new symptoms, even if mild, self-isolate and arrange a COVID-19 test by either phoning your GP, booking online or going to a test centre.

<u>Self-isolation – (stay in your room).</u> Self-isolation means staying indoors in a room on your own if possible, and completely avoiding contact with other people, including others in your household. Self-isolation is for people who have had a Reverse Transcription Polymerase Chain Reaction (RT-PCR) test that shows they have COVID-19 and also for people who are thought to have COVID-19, but are still waiting for a test or test result.

<u>Restricting movement</u> – (stay at home). Restricting movements means staying at home and avoiding contact with other people and social situations as much as possible. Restricting movement is for people who do not have signs of COVID-19 but who may be at higher risk of getting COVID-19 because they were exposed to a particular risk.

These guidelines do not include information for those who have recently returned from a designated state. Please see <u>gov.ie</u> for information for those returning from designated states.

If you are a health care worker, please refer to occupational health guidelines.

Please see pages 2 - 4 for full guide.

	Situation	Actions for person	Actions for others
	Situation	Actions for person	living with or in close
1	Positive COVID-19 RT-PCR test	You need to self-isolate for 10 full days from when the symptoms of COVID-19 started, with no fever for the last 5 days. If you do not have symptoms of COVID-19, you must self-isolate for 10 full days, starting from the date of the positive test, even if you are vaccinated .	contact with person Close contacts must restrict their movements for 14 days after last contact with case ¹ . Close contacts will be tested on Day 0 and Day 10 after last contact with the case. If the test result on Day 10 is "Negative/Not Detected" and the close contact does not have any symptoms of COVID-19, they can stop restricting their movements.
2	A person with COVID-19 symptoms awaiting a COVID-19 RT-PCR test or test result. Sometimes the GP may say that results need to be repeated to be sure if the person has COVID-19 or not. This section applies for those in this situation also.	You need to self-isolate, including from other people in the house if possible, while waiting for the COVID-19 test or test result, even if you are vaccinated.	Household contacts ² must restrict their movements while waiting for the COVID-19 test result, or until the person has been told that their illness is not from COVID-19.
3	A person with symptoms of COVID-19 disease and a "Negative/Not Detected" COVID-19 test result	You should follow medical advice from your GP. Self-isolate at home until 48 hours after your symptoms have resolved. Continue to limit contact with other people that you live with while you have symptoms.	There is no need for close contacts or household contacts ² to restrict their movements any more.
4	A person with symptoms of illness that may be COVID-19, whose symptoms may also relate to an underlying medical condition, who is waiting for a doctor to assess their illness and decide if a COVID-19 test is needed.	You need to self-isolate until you have been assessed by a doctor, even if you are vaccinated. If the doctor thinks that you need to take a COVID-19 test based on your symptoms, follow advice in row 2 above.	Other people do not need to restrict their movements until the person has been assessed by a doctor. If the doctor decides the person needs a COVID- 19 test, household contacts ² of the person need to restrict their movements while waiting for the COVID-19 test result (Follow advice in Row 2).
5	A person with no symptoms of COVID-19 disease, but they have been told they are a close contact of someone with COVID-19. ³	You need to restrict your movements for 14 days. ¹ You will be tested on Day 0 and Day 10 after your last contact with the case. If the test result on Day 10 is "Negative/Not Detected" and you do not have any symptoms of COVID-19, you can stop restricting your movements.	No restrictions are required. No testing of household contacts ² required.
6	A person with no symptoms of COVID-19 disease, but they have been told that they are a close contact of someone with COVID-19, however they have had COVID-19 infection themselves in the past 9 months.	 If you received a positive result for COVID-19 in the last 9 months and you now have no symptoms, you do not need to be tested or to restrict your movements. If it is more than 9 months since you received a positive result for COVID- 19 and you have been told that you are a close contact with a COVID-19 case, but have no symptoms, you 	No restrictions are required, unless the person develops symptoms.

		need to restrict your movements and	
		 be tested (follow advice in row 5). If you start to develop symptoms you 	
		will need to self-isolate and be	
		tested for COVID-19, even if it is less	
		than 9 months since you were	
		diagnosed with COVID-19. (follow	
		advice in row 2)	
7	A person with no symptoms of COVID-19	Persons who are fully vaccinated, include those who are:	No restrictions are
-	disease, but they have been told that they are a close contact of someone	a) 7 days after receipt of the second	required, unless the person develops
	with COVID-19 ⁴ , however they have	Pfizer-BioNTech (Comirnaty®) dose	symptoms.
	been fully vaccinated.	(two dose vaccination course)	, ,
		b) 14 days after receipt of the second	
		Moderna (Moderna [®]) dose (two	
		dose vaccination course)	
		 c) 14 days after receipt of the Janssen (Janssen[®]) dose (one dose 	
		vaccination course)	
		d) 15 days after receipt of the second	
		AstraZeneca (Vaxzevria® or	
		Covishield) dose (two dose	
		vaccination course)	
		 If you have been fully vaccinated and remain asymptomatic, you do not 	
		need to be tested or to restrict your	
		movements unless specific	
		circumstances apply ⁴ .	
		If you start to develop symptoms you	
		will need to self-isolate and be	
		tested for COVID-19, even if you	
		have been fully vaccinated (follow advice in row 2).	
0	A person with COVID-19 symptoms	You need self-isolate and be tested for	Household contacts ²
8	AND	COVID-19. If your test result is negative,	must restrict movements
	the person is also a close contact of a	remain at home and self-isolate until 48 hours	while waiting for the
	confirmed case of COVID 19	after resolution of all symptoms. After this	COVID-19 test result.
		you must continue to restrict your movements for 14 days from the day of last	
		contact with the case and be tested at day 0	
		and day 10. (Follow advice in row 5)	
9	A person with COVID-19 symptoms	You need self-isolate and be tested for	Household contacts ²
9	AND	COVID-19. If your COVID-19 test is positive,	must restrict movements
	the person has had NO contact with a	follow advice in row 1. If your COVID-19 test	while waiting for the
	COVID-19 case A person who has travelled into Ireland	is negative, follow advice in Row 3. If you have valid proof of vaccination, no	COVID-19 test result. No restrictions unless
10	from a <u>non-</u> designated state.	travel-related testing or quarantine ⁵ will be	the person who has
		necessary.	returned from abroad
		If you have valid proof of recovery from	develops symptoms.
		COVID in the past 180 days, no travel-related	
		testing or quarantine will be necessary.	
		If you do not have valid proof of vaccination or recovery, you will need to:	
		 present evidence of a negative result 	
		from a RT-PCR test taken within 72	
		hours prior to arrival into the	
		country	
		 <u>self-quarantine</u> for 14 days 	

		 if you receive a negative result from a RT-PCR test taken from day 5 onwards after arrival into Ireland, you will be able to leave quarantine 	
11	A person who has COVID-19 symptoms, fits the criteria to be tested for COVID 19, but not tested for any reason	You need to self-isolate for 10 full days from when the symptoms of COVID-19 started with no fever for the last 5 days	If close contacts are symptomatic, you will need to restrict movements and be tested, even if vaccinated. If asymptomatic, please follow advice in row 5.
12	A person who is a confirmed case of COVID-19 and is discharged from hospital to a residential care facility (RCF) before they have completed their 14-day period of self-isolation	The person must complete the remainder of the 14-day period of self-isolation in the RCF, with the last 5 days fever free ⁶	Carers must use transmission-based precautions
13	A person who had not been diagnosed in the last 14 days with COVID-19 is discharged from hospital to a residential care facility (RCF)	The person must restrict their movements for 14 days after they arrive at the RCF, unless they are fully vaccinated or have had COVID 19 infection in the past 9 months, in which case they are not required to restrict movements ⁶	Carers can use standard precautions

- In limited cases it might not be possible for a case to self-isolate from their household contacts. This may occur
 if the case has additional needs. In these exceptional circumstances, household contacts must restrict their
 movements for <u>17 days from the day of onset of symptoms in the case (or test date if asymptomatic). This is
 regardless of whether further cases are identified in the household.
 </u>
- 2. Household contacts are people living or sleeping in the same house, people in shared living spaces who share a kitchen or bathroom, and sexual partners.
- 3. If you are a close contact with a case of COVID-19, but you are fully vaccinated, please refer to '<u>Guidance on the impact of vaccination on contact tracing</u>' Known contact with a case of COVID-19 in which the case is a Person Under Investigation, probable or confirmed variant of concern excluding Alpha and Delta. In this situation the close contact should be managed as a close contact of a VOC.
- 4. If the person's immune system response to vaccination could be compromised due to either a known medical condition or being on immunosuppressive treatment, they should be treated as a close contact offered two tests and advised to restrict their movements. If there is any uncertainty as to whether the close contact has a medical condition or takes a treatment that would result in a sub-optimal response to vaccination, they should also be advised to restrict their movements and contact their treating physician who can advise if these recommendations apply to them. See Appendix B for a list of medical conditions and immunosuppressive treatments which are associated with sub-optimal response to vaccines. This list was compiled from 'The Immunisation Guidelines for Ireland', Chapter 5a COVID-19 and may be subject to change/update in future.
 - A public health or occupational health risk assessment has identified other specific grounds for concern
 - The close contact develops symptoms of COVID-19, in which case they need to immediately self-isolate and be referred for one test. If the test result is negative they can discontinue self-isolation once they are symptom free for 48 hours.
- 5. Quarantine is a legal term which is used by the government to describe the obligation for persons who have returned to Ireland, to either (a) stay at home and self-isolate from others if they have returned from a non-designated state, or (b) to isolate in a mandatory quarantine hotel if they have returned from a designated state.

6. See <u>'Interim Public Health, Infection Prevention & Control Guidelines on the Prevention and Management</u> of COVID-19 Cases and Outbreaks in Residential Care Facilities', for further information